

A CATEGORICAL APPROACH TO SYNTACTIC MONOIDS

JIŘÍ ADÁMEK^a, STEFAN MILIUS^b, AND HENNING URBAT^c

^a Institut für Theoretische Informatik, Technische Universität Braunschweig, Germany
e-mail address: adamek@iti.cs.tu-bs.de

^{b,c} Lehrstuhl für Theoretische Informatik, Friedrich-Alexander-Universität Erlangen-Nürnberg, Germany
e-mail address: mail@stefan-milius.eu, henning.urbat@fau.de

ABSTRACT. The syntactic monoid of a language is generalized to the level of a symmetric monoidal closed category \mathcal{D} . This allows for a uniform treatment of several notions of syntactic algebras known in the literature, including the syntactic monoids of Rabin and Scott ($\mathcal{D} = \text{sets}$), the syntactic ordered monoids of Pin ($\mathcal{D} = \text{posets}$), the syntactic semirings of Polák ($\mathcal{D} = \text{semilattices}$), and the syntactic associative algebras of Reutenauer ($\mathcal{D} = \text{vector spaces}$). Assuming that \mathcal{D} is a commutative variety of algebras or ordered algebras, we prove that the syntactic \mathcal{D} -monoid of a language L can be constructed as a quotient of a free \mathcal{D} -monoid modulo the syntactic congruence of L , and that it is isomorphic to the transition \mathcal{D} -monoid of the minimal automaton for L in \mathcal{D} . Furthermore, in the case where the variety \mathcal{D} is locally finite, we characterize the regular languages as precisely the languages with finite syntactic \mathcal{D} -monoids.

1. INTRODUCTION

One of the successes of the theory of coalgebras is that ideas from automata theory can be developed at a level of abstraction where they apply uniformly to many different types of systems. In fact, classical deterministic automata are a standard example of coalgebras for an endofunctor. And that automata theory can be studied with coalgebraic methods rests on the observation that formal languages form the final coalgebra.

The present paper contributes to a new category-theoretic view of *algebraic* automata theory. In this theory one starts with an elegant machine-independent notion of language recognition: a language $L \subseteq X^*$ is *recognized* by a monoid morphism $e : X^* \rightarrow M$ if it is the preimage under e of some subset of M . Regular languages are then characterized as precisely the languages recognized by finite monoids. A key concept, introduced by

Key words and phrases: Syntactic monoid, transition monoid, algebraic automata theory, duality, coalgebra, algebra, symmetric monoidal closed category, commutative variety.

Stefan Milius acknowledges support by the Deutsche Forschungsgemeinschaft (DFG) under project MI 717/5-1.

Jiří Adámek and Henning Urbat acknowledge support by the Deutsche Forschungsgemeinschaft (DFG) under project AD 187/2-1.

Rabin and Scott [19] (and earlier in unpublished work of Myhill), is the *syntactic monoid* of a language L . It serves as a canonical algebraic recognizer of L , namely the smallest X -generated monoid recognizing L . Two standard methods to construct the syntactic monoid are:

- (1) as a quotient of the free monoid X^* modulo the *syntactic congruence* of L , which is a two-sided version of the well-known Myhill-Nerode equivalence, and
- (2) as the *transition monoid* of the minimal automaton for L .

In addition to syntactic monoids there are several related notions of syntactic algebras for (weighted) languages in the literature, most prominently the syntactic ordered monoids of Pin [16], the syntactic idempotent semirings of Polák [18] and the syntactic associative algebras of Reutenauer [20], all of which admit constructions similar to (1) and (2). A crucial observation is that monoids, ordered monoids, idempotent semirings and associative algebras are precisely the monoid objects in the categories of sets, posets, semilattices and vector spaces, respectively. Moreover, all these categories are symmetric monoidal closed w.r.t. their usual tensor product.

The main goal of our paper is to develop a theory of algebraic recognition in a general symmetric monoidal closed category $\mathcal{D} = (\mathcal{D}, \otimes, I)$, covering all the above syntactic algebras uniformly. Following Goguen [12], a *language* in \mathcal{D} is a morphism $L : X^{\otimes} \rightarrow Y$ where X is a fixed object of inputs, Y is a fixed object of outputs, and X^{\otimes} denotes the free \mathcal{D} -monoid on X . And a \mathcal{D} -*automaton* is given by the picture below: it consists of an object of states Q , a morphism i representing the initial state, an output morphism f , and a transition morphism δ which may be presented in its curried form $\lambda\delta$.

$$\begin{array}{ccc}
 & X \otimes Q & \\
 & \downarrow \delta & \\
 I & \xrightarrow{i} Q \xrightarrow{f} Y & \\
 & \downarrow \lambda\delta & \\
 & [X, Q] &
 \end{array} \tag{1.1}$$

As observed by Rutten [21], this means that an automaton is at the same time an *algebra* $I + X \otimes Q \xrightarrow{[i, \delta]} Q$ for the functor $FQ = I + X \otimes Q$, and a *coalgebra* $Q \xrightarrow{\langle f, \lambda\delta \rangle} Y \times [X, Q]$ for the functor $TQ = Y \times [X, Q]$. It turns out that much of the classical (co-)algebraic theory of automata in the category of sets extends to this level of generality. Thus Goguen [12] demonstrated that the initial algebra for F coincides with the free \mathcal{D} -monoid X^{\otimes} , and that every language is accepted by a unique minimal \mathcal{D} -automaton. We will add to this picture the observation that the final coalgebra for T is carried by the object of languages $[X^{\otimes}, Y]$, see Proposition 2.26.

In Section 3 we introduce the central concept of our paper, the *syntactic \mathcal{D} -monoid* of a language $L : X^{\otimes} \rightarrow Y$, which by definition is the smallest X -generated \mathcal{D} -monoid recognizing L . In Corollary 3.10 we give a natural condition on a monoidal category that ensures the existence of a syntactic \mathcal{D} -monoid for every language L . Furthermore, assuming that \mathcal{D} is a commutative variety of algebras or ordered algebras, we will show that the above constructions (1) and (2) for the classical syntactic monoid adapt to our general setting: the syntactic \mathcal{D} -monoid is (1) the quotient of X^{\otimes} modulo the syntactic congruence of L (Theorem 3.14), and (2) the transition \mathcal{D} -monoid of the minimal \mathcal{D} -automaton for L (Theorem 4.7). As

special instances we recover syntactic monoids ($\mathcal{D} = \text{sets}$), syntactic ordered monoids ($\mathcal{D} = \text{posets}$), syntactic semirings ($\mathcal{D} = \text{semilattices}$) and syntactic associative algebras ($\mathcal{D} = \text{vector spaces}$). In addition, our categorical setting yields new types of syntactic algebras “for free”. For example, we will identify monoids with zero as the algebraic structures representing partial automata (the case $\mathcal{D} = \text{pointed sets}$) and obtain the *syntactic monoid with zero* for a given language. Similarly, by taking as \mathcal{D} the variety of algebras with an involutive unary operation we obtain *syntactic involution monoids*.

Most of the results of our paper apply to arbitrary languages. In Section 5 we will investigate *\mathcal{D} -regular languages*, that is, languages accepted by \mathcal{D} -automata with a finitely presentable object of states. Under suitable assumptions on \mathcal{D} , we will prove that a language is \mathcal{D} -regular iff its syntactic \mathcal{D} -monoid is carried by a finitely presentable object (Theorem 5.6). We will also derive a dual characterization of the syntactic \mathcal{D} -monoid which is new even in the “classical” case $\mathcal{D} = \text{sets}$: if \mathcal{D} is a locally finite variety, i.e. all finitely generated algebras are finite, and if moreover some other locally finite variety \mathcal{C} is dual to \mathcal{D} on the level of finite objects, the syntactic \mathcal{D} -monoid of L dualizes to the local variety of languages in \mathcal{C} generated by the reversed language of L .

Related work. This paper is a reworked full version of the conference paper [2]. Apart from including full proofs, it has two new contributions. First, we present in Section 3.1 a purely categorical existence criterion for syntactic monoids in an abstract symmetric monoidal closed category, whereas in [2] only a set-theoretic construction was given. Secondly, we treat the case of ordered algebras explicitly, covering new examples such as the syntactic ordered monoids of Pin [16].

Another categorical approach to (classical) syntactic monoids appears in the work of Ballester-Bolínches, Cosme-López and Rutten [4]. These authors consider automata in the category of sets specified by *equations* or dually by *coequations*, which leads to a construction of the automaton underlying the syntactic monoid of a language. The fact that it forms the transition monoid of a minimal automaton is also interpreted in that setting. In the present paper we take a more general approach by studying algebraic recognition in an arbitrary symmetric monoidal closed category \mathcal{D} . One important source of inspiration for our categorical setting was the work of Goguen [12].

In the recent papers [1, 3] we presented a categorical view of *varieties of languages*, another central topic of algebraic automata theory. Building on the duality-based approach of Gehrke, Grigorieff and Pin [11], we generalized Eilenberg’s variety theorem and its local version to the level of an abstract (pre-)duality between algebraic categories. The idea to replace monoids by monoid objects in a commutative variety \mathcal{D} originates in this work.

Another categorical setting for algebraic language theory can be found in Bojányk [8]. He considers, in lieu of monoids in commutative varieties, Eilenberg-Moore algebras for an arbitrary monad on sorted sets, and defines syntactic congruences in this more general setting. Our Theorem 3.14 is a special case of [8, Theorem 3.1].

2. PRELIMINARIES

Throughout this paper we investigate deterministic automata in a symmetric monoidal closed category. In all concrete applications, this category will be a commutative variety of algebras or ordered algebras. Recall that for a finitary signature Σ a *variety of Σ -algebras* is a class of Σ -algebras specified by equations $s = t$ between Σ -terms. Equivalently, by Birkhoff’s HSP theorem, it is a class of Σ -algebras closed under products, subalgebras and

quotients (= homomorphic images). Similarly, *ordered Σ -algebras* are posets equipped with order-preserving Σ -operations, and their morphisms are order-preserving Σ -homomorphisms. A *quotient* of an ordered algebra B is represented by a surjective morphism $e: B \twoheadrightarrow A$, and a *subalgebra* of B is represented by an *order-reflecting* morphism $m: A \twoheadrightarrow B$, i.e. $mx \leq my$ iff $x \leq y$. A *variety of ordered Σ -algebras* is a class of ordered Σ -algebras specified by inequations $s \leq t$ between Σ -terms. Equivalently, by Bloom's HSP theorem [7], it is a class of ordered Σ -algebras closed under products, subalgebras and quotients.

A variety \mathcal{D} of algebras or ordered algebras is called *commutative* (a.k.a. *entropic*) if, for any two algebras A and B in \mathcal{D} , the set $[A, B]$ of all homomorphisms from A to B forms a subalgebra $[A, B] \twoheadrightarrow B^{|A|}$ of the product of $|A|$ copies of B . Equivalently, the corresponding monad over **Set** (resp. **Pos**, the category of posets) is commutative in the sense of Kock [13].

Example 2.1. In our applications, we shall work with the following commutative varieties:

- (1) **Set** is a commutative variety (over the empty signature) with $[A, B] = B^A$.
- (2) **Pos** is a commutative variety of ordered algebras (over the empty signature). Here $[A, B]$ is the poset of monotone functions from A to B , ordered pointwise.
- (3) A *pointed set* (A, \perp) is a set A together with a chosen point $\perp \in A$. The category **Set** $_{\perp}$ of pointed sets and point-preserving functions is a commutative variety. The point of $[(A, \perp_A), (B, \perp_B)]$ is the constant function with value \perp_B .
- (4) An *involution algebra* is a set with an involutive unary operation $x \mapsto \tilde{x}$, i.e. $\tilde{\tilde{x}} = x$. We call \tilde{x} the *complement* of x . Morphisms are functions f preserving complements, i.e. $f(\tilde{x}) = \widetilde{f(x)}$. The variety **Inv** of involution algebras is commutative. Indeed, the set $[A, B]$ of all homomorphisms is an involution algebra with pointwise complementation.
- (5) All other examples we treat in our paper are varieties of modules over a semiring. Recall that a *semiring* $\mathbb{S} = (S, +, \cdot, 0, 1)$ consists of a commutative monoid $(S, +, 0)$ and a monoid $(S, \cdot, 1)$ subject the following equational laws:

$$0s = s0 = 0, \quad r(s + t) = rs + rt \quad \text{and} \quad (r + s)t = rt + st.$$

A *module* over a semiring \mathbb{S} is a commutative monoid $(M, +, 0)$ together with a scalar multiplication $\cdot : S \times M \rightarrow M$ (written simply as juxtaposition $(s, x) \mapsto sx$, as usual) such that the following laws hold:

$$\begin{aligned} (r + s)x &= rx + sx, & r(x + y) &= rx + ry, & (rs)x &= r(sx), \\ 0x &= 0, & 1x &= x, & r0 &= 0. \end{aligned}$$

We denote by **Mod**(\mathbb{S}) the category of \mathbb{S} -modules and module homomorphisms (i.e. \mathbb{S} -linear maps). This is a commutative variety; here $[A, B]$ is the set of all module homomorphisms with pointwise module structure. Three interesting special cases of **Mod**(\mathbb{S}) are:

- (a) $\mathbb{S} = \{0, 1\}$, the boolean semiring with $1 + 1 = 1$: the category **JSL** $_0$ of join-semilattices with the least element 0, and homomorphisms preserving joins and 0;
- (b) $\mathbb{S} = \mathbb{Z}$: the category **Ab** of abelian groups and group homomorphisms;
- (c) $\mathbb{S} = \mathbb{K}$ (a field): the category **Vec**(\mathbb{K}) of vector spaces over \mathbb{K} and linear maps.

Notation 2.2. For any variety \mathcal{D} of algebras or ordered algebras, we denote by $\Psi : \mathbf{Set} \rightarrow \mathcal{D}$ the left adjoint to the forgetful functor $|-| : \mathcal{D} \rightarrow \mathbf{Set}$. Thus ΨX_0 is the free object of \mathcal{D} on the set X_0 .

- Example 2.3.** (1) For $\mathcal{D} = \mathbf{Set}$ or \mathbf{Pos} we have $\Psi X_0 = X_0$ (discretely ordered).
(2) For $\mathcal{D} = \mathbf{Set}_\perp$ the free pointed set on X_0 is $\Psi X_0 = X_0 + \{\perp\}$.
(3) For $\mathcal{D} = \mathbf{Inv}$ the free involution algebra on X_0 is $\Psi X_0 = X_0 + \widetilde{X}_0$ where \widetilde{X}_0 is a copy of X_0 (whose elements are denoted \tilde{x} for $x \in X_0$). The involution swaps the copies of X_0 , and the universal arrow $X_0 \rightarrow X_0 + \widetilde{X}_0$ is the left coproduct injection.
(4) For $\mathcal{D} = \mathbf{Mod}(\mathbb{S})$ the free module ΨX_0 is the submodule of \mathbb{S}^{X_0} on all functions $X_0 \rightarrow \mathbb{S}$ with finite support. Equivalently, ΨX_0 consists of formal linear combinations $\sum_{i=1}^n s_i x_i$ with $s_i \in \mathbb{S}$ and $x_i \in X_0$. In particular, $\Psi X_0 = \mathcal{P}_f X_0$ (finite subsets of X_0) for $\mathcal{D} = \mathbf{JSL}_0$, and ΨX_0 is the vector space with basis X_0 for $\mathcal{D} = \mathbf{Vec}(\mathbb{K})$.

Definition 2.4. Given objects A, B and C of a variety \mathcal{D} , a *bimorphism* from A, B to C is a function $f : |A| \times |B| \rightarrow |C|$ such that the maps $f(a, -) : |B| \rightarrow |C|$ and $f(-, b) : |A| \rightarrow |C|$ carry morphisms of \mathcal{D} for every $a \in |A|$ and $b \in |B|$. A *tensor product* of A and B is a universal bimorphism $\eta_{A,B} : |A| \times |B| \rightarrow |A \otimes B|$, which means that for every bimorphism $f : |A| \times |B| \rightarrow |C|$ there is a unique morphism $f' : A \otimes B \rightarrow C$ in \mathcal{D} with $f' \cdot \eta_{A,B} = f$.

Theorem 2.5 (Banaschewski and Nelson [5]). *Every commutative variety \mathcal{D} of algebras or ordered algebras has tensor products, making $(\mathcal{D}, \otimes, I)$ with $I = \Psi 1$ a symmetric monoidal closed category. That is, we have the following bijective correspondence of morphisms, natural in $A, B, C \in \mathcal{D}$:*

$$\frac{f : A \otimes B \rightarrow C}{\lambda f : A \rightarrow [B, C]}$$

Remark 2.6. We will use the following properties of the tensor product, see [5]:

- (1) $- \otimes B$ and $A \otimes -$ are left-adjoints and hence preserve all colimits.
- (2) The morphism λf is obtained by currying the bimorphism $f' = f \cdot \eta_{A,B} : |A| \times |B| \rightarrow |C|$ corresponding to f , that is, $\lambda f(a)(b) = f'(a, b)$ for $a \in |A|$ and $b \in |B|$.
- (3) The evaluation morphism $\text{ev} = \text{ev}_{A,B} : [A, B] \otimes A \rightarrow B$, i.e. the counit of the adjunction $- \otimes A \dashv [A, -]$, is the morphism in \mathcal{D} corresponding to the bimorphism

$$\text{ev}' : [|A, B|] \times |A| \rightarrow |B|, \quad (h, a) \mapsto h(a).$$

- (4) The right unit isomorphism $i : A \cong A \otimes I$ is given by

$$i = (|A| \xrightarrow{\langle A, c_1 \rangle} |A| \times |I| \xrightarrow{\eta} |A \otimes I|)$$

where $c_1 : |A| \rightarrow |I|$ is the constant map that sends $a \in A$ to the generator of $I = \Psi 1$.

Assumptions 2.7. For the rest of this paper let \mathcal{D} be a commutative variety of algebras or ordered algebras, equipped with the monoidal structure $(\mathcal{D}, \otimes, I)$ of the above theorem. Furthermore, we fix an object X (of inputs) and an object Y (of outputs) in \mathcal{D} .

Remark 2.8. Despite the above algebraic assumptions we will state many of our definitions, theorems and proofs for an arbitrary symmetric monoidal closed category $(\mathcal{D}, \otimes, I)$ subject to additional properties.

Remark 2.9. Recall that a *monoid* (M, m, i) in a monoidal category $(\mathcal{D}, \otimes, I)$ (with tensor product $\otimes : \mathcal{D} \times \mathcal{D} \rightarrow \mathcal{D}$ and tensor unit $I \in \mathcal{D}$) is an object M equipped with a multiplication $m : M \otimes M \rightarrow M$ and unit $i : I \rightarrow M$ satisfying the usual associative and unit laws. Due to \otimes and $I = \Psi 1$ representing bimorphisms, this categorical definition is equivalent to the following algebraic one in our setting: a \mathcal{D} -*monoid* is a triple (M, \bullet, i) where M is an object of \mathcal{D} and $(|M|, \bullet, i)$ is a monoid (in the usual sense) with $\bullet : |M| \times |M| \rightarrow |M|$ a bimorphism

of \mathcal{D} . A *morphism* $h : (M, \bullet, i) \rightarrow (M', \bullet', i')$ of \mathcal{D} -monoids is a morphism $h : M \rightarrow M'$ of \mathcal{D} such that $|h| : |M| \rightarrow |M'|$ is a monoid morphism. We denote by $\mathbf{Mon}(\mathcal{D})$ the category of \mathcal{D} -monoids and their homomorphisms. In the following we will freely work with \mathcal{D} -monoids in both categorical and algebraic disguise.

- Example 2.10.** (1) In \mathbf{Set} the tensor product is the cartesian product, $I = \{*\}$, and \mathbf{Set} -monoids are ordinary monoids.
- (2) In \mathbf{Pos} the tensor product is the cartesian product, $I = \{*\}$, and \mathbf{Pos} -monoids are *ordered monoids*, that is, posets equipped with a monoid structure whose multiplication \bullet is monotone.
- (3) In \mathbf{Set}_\perp we have $I = \{\perp, *\}$, and the tensor product of pointed sets (A, \perp_A) and (B, \perp_B) is $A \otimes B = (A \setminus \{\perp_A\}) \times (B \setminus \{\perp_B\}) + \{\perp\}$. \mathbf{Set}_\perp -monoids are precisely monoids with zero. Indeed, given a \mathbf{Set}_\perp -monoid structure on (A, \perp) we have $x \bullet \perp = \perp = \perp \bullet x$ for all x because \bullet is a bimorphism, i.e. \perp is a zero element. Morphisms of $\mathbf{Mon}(\mathbf{Set}_\perp)$ are zero-preserving monoid morphisms.
- (4) An \mathbf{Inv} -monoid (also called an *involution monoid*) is a monoid equipped with an involution $x \mapsto \tilde{x}$ such that $x \bullet \tilde{y} = \tilde{x} \bullet y = \widetilde{x \bullet y}$. For example, for any set A the power set $\mathcal{P}A$ naturally carries the structure of an involution monoid: the involution takes complements, $\tilde{S} = A \setminus S$, and the monoid multiplication is the symmetric difference $S \oplus T = (S \setminus T) \cup (T \setminus S)$.
- (5) \mathbf{JSL}_0 -monoids are precisely idempotent semirings (with 0 and 1). Indeed, a \mathbf{JSL}_0 -monoid on a semilattice (i.e. a commutative idempotent monoid) $(D, +, 0)$ is given by a unit 1 and a monoid multiplication that, being a bimorphism, distributes over $+$ and 0.
- (6) More generally, a $\mathbf{Mod}(\mathbb{S})$ -monoid is precisely an associative algebra over \mathbb{S} : it consists of an \mathbb{S} -module together with a unit 1 and a monoid multiplication that distributes over $+$ and 0 and moreover preserves scalar multiplication in both components.

Notation 2.11. We denote by $X^{\otimes n}$ ($n < \omega$) the n -fold tensor power of X , recursively defined by $X^{\otimes 0} = I$ and $X^{\otimes(n+1)} = X \otimes X^{\otimes n}$.

Proposition 2.12 (see Mac Lane [14]). *Let \mathcal{D} be a symmetric monoidal closed category with countable coproducts. Then the forgetful functor $\mathbf{Mon}(\mathcal{D}) \rightarrow \mathcal{D}$ has a left adjoint assigning to every object X the free \mathcal{D} -monoid $X^\otimes = \coprod_{n < \omega} X^{\otimes n}$. The monoid structure (X^\otimes, m_X, i_X) is given by the coproduct injection $i_X : I = X^{\otimes 0} \rightarrow X^\otimes$ and $m_X : X^\otimes \otimes X^\otimes \rightarrow X^\otimes$, where $X^\otimes \otimes X^\otimes \cong \coprod_{n, k < \omega} X^{\otimes n} \otimes X^{\otimes k}$ and m_X has as its (n, k) -component the $(n+k)$ -th coproduct injection. The universal arrow $\eta_X : X \rightarrow X^\otimes$ is the first coproduct injection.*

Remark 2.13. For any \mathcal{D} -monoid (M, m, i) and any morphism $f : X \rightarrow M$ the extension to the unique \mathcal{D} -monoid morphism $f^+ : X^\otimes \rightarrow M$ with $f^+ \cdot \eta_X = f$ is defined inductively on the components of the coproduct $\coprod_{n < \omega} X^{\otimes n}$ as follows: $f^+ = [f_n^+]_{n < \omega}$, where $f_0^+ = i : X^{\otimes 0} = I \rightarrow M$ and

$$f_{n+1}^+ = (X^{\otimes(n+1)} = X \otimes X^{\otimes n} \xrightarrow{f \otimes f_n^+} M \otimes M \xrightarrow{m} M).$$

In our setting where \mathcal{D} is commutative variety we have the following construction of free \mathcal{D} -monoids on free \mathcal{D} -objects:

Proposition 2.14. *The free \mathcal{D} -monoid on $X = \Psi X_0$ is $X^\otimes = \Psi X_0^*$. Its monoid multiplication extends the concatenation of words in X_0^* , and its unit is the empty word ε .*

Proof. A constructive proof can be found in [1]. Here we sketch a more conceptual argument, using the universal property of the tensor product. Observe that the free-algebra functor $\Psi : \mathbf{Set} \rightarrow \mathcal{D}$ is strongly monoidal, i.e. it preserves the unit and tensor product up to natural isomorphism. Indeed, we have $\Psi 1 = I$ by definition. To see that $\Psi(A \times B) \cong \Psi A \otimes \Psi B$ for all sets A and B , consider the following bijections (natural in $D \in \mathcal{D}$):

$$\begin{aligned} \mathcal{D}(\Psi(A \times B), D) &\cong \mathbf{Set}(A \times B, |D|) \\ &\cong \mathbf{Set}(A, |D|^B) \\ &\cong \mathbf{Set}(A, |[\Psi B, D]|) \\ &\cong \mathcal{D}(\Psi A, [\Psi B, D]) \\ &\cong \mathcal{D}(\Psi A \otimes \Psi B, D). \end{aligned}$$

This implies $\Psi(A \times B) \cong \Psi A \otimes \Psi B$ by the Yoneda lemma. Using the fact that Ψ preserves coproducts, being a left adjoint, we conclude

$$X^\otimes \cong \coprod_{n < \omega} X^{\otimes n} \cong \coprod_{n < \omega} \Psi X_0^n \cong \Psi\left(\coprod_{n < \omega} X_0^n\right) = \Psi X_0^*. \quad \square$$

Example 2.15. (1) In \mathbf{Set} and \mathbf{Pos} we have $X^\otimes = X^*$ (discretely ordered).

(2) In \mathbf{Set}_\perp with $X = \Psi X_0 = X_0 + \{\perp\}$ we get $X^\otimes = X_0^* + \{\perp\}$. The product $x \bullet y$ is concatenation for $x, y \in X_0^*$, and otherwise \perp .

(3) In \mathbf{Inv} with $X = \Psi X_0 = X_0 + \widetilde{X}_0$ we have $X^\otimes = X_0^* + \widetilde{X}_0^*$. The multiplication restricted to X_0^* is concatenation, and is otherwise determined by $\widetilde{u} \bullet v = \widetilde{u}v = u \bullet \widetilde{v}$ for $u, v \in X_0^*$.

(4) In \mathbf{JSL}_0 with $X = \Psi X_0 = \mathcal{P}_f X_0$ we have $X^\otimes = \mathcal{P}_f X_0^*$, the semiring of all finite languages over X_0 . Its addition is union and its multiplication is the concatenation of languages.

(5) More generally, in $\mathbf{Mod}(\mathbb{S})$ with $X = \Psi X_0$ we get $X^\otimes = \Psi X_0^* = \mathbb{S}[X_0]$, the module of all polynomials over \mathbb{S} in finitely many variables from X_0 , i.e., $\sum_{i=1}^n c(w_i)w_i$ with $w_i \in X_0^*$ and $c(w_i) \in \mathbb{S}$. Hence the elements of $\mathbb{S}[X_0]$ are functions $c : X_0^* \rightarrow \mathbb{S}$ with finite support, i.e. finite weighted languages. The \mathbb{S} -algebraic structure of $\mathbb{S}[X_0]$ is given by the usual addition, scalar multiplication and product of polynomials.

Definition 2.16 (Goguen [12]). A \mathcal{D} -automaton (Q, δ, i, f) consists of an object Q (of states) and morphisms $\delta : X \otimes Q \rightarrow Q$, $i : I \rightarrow Q$ and $f : Q \rightarrow Y$; see Diagram (1.1). An automata homomorphism $h : (Q, \delta, i, f) \rightarrow (Q', \delta', i', f')$ is a morphism $h : Q \rightarrow Q'$ preserving transitions as well as initial states and outputs, i.e. making the following diagrams commute:

$$\begin{array}{ccc} X \otimes Q & \xrightarrow{\delta} & Q \\ X \otimes h \downarrow & & \downarrow h \\ X \otimes Q' & \xrightarrow{\delta'} & Q' \end{array} \quad \begin{array}{ccccc} I & \xrightarrow{i} & Q & \xrightarrow{f} & Y \\ & \searrow i' & \downarrow h & \searrow f' & \\ & & Q' & & \end{array}$$

The above definition makes sense in any monoidal category \mathcal{D} . In our algebraic setting, since $I = \Psi 1$, the morphism i chooses an *initial state* in $|Q|$. Moreover, if $X = \Psi X_0$ for some set X_0 (of inputs), the morphism δ amounts to a choice of endomorphisms $\delta_a : Q \rightarrow Q$ for $a \in X_0$, representing transitions. This follows from the bijections

$$\frac{\Psi X_0 \otimes Q \rightarrow Q \quad \text{in } \mathcal{D}}{\Psi X_0 \rightarrow [Q, Q] \quad \text{in } \mathcal{D}} \quad \frac{\quad}{X_0 \rightarrow \mathcal{D}(Q, Q) \quad \text{in } \mathbf{Set}}$$

We will occasionally write $q \xrightarrow{a} q'$ for $\delta_a(q) = q'$ if δ is clear from the context.

- Example 2.17.** (1) The classical deterministic automata are the case $\mathcal{D} = \mathbf{Set}$ and $Y = \{0, 1\}$. Here $f : Q \rightarrow \{0, 1\}$ defines the set $F = f^{-1}[1] \subseteq Q$ of final states. For a general set Y we get deterministic Moore automata with outputs in Y .
- (2) For $\mathcal{D} = \mathbf{Pos}$ and $Y = \{0 < 1\}$ we get *ordered* deterministic automata. That is, automata with a partially ordered set Q of states, monotone transition maps δ_a , and an upper set $F = f^{-1}[1] \subseteq Q$ of final states.
- (3) The setting $\mathcal{D} = \mathbf{Set}_\perp$ with $X = X_0 + \{\perp\}$ and $Y = \{\perp, 1\}$ gives *partial* deterministic automata. Indeed, the state object (Q, \perp) has transitions $\delta_a : (Q, \perp) \rightarrow (Q, \perp)$ for $a \in X_0$ preserving \perp , that is, \perp is a sink state. Equivalently, we may consider δ_a as a partial transition map on the state set $Q \setminus \{\perp\}$. The morphism $f : (Q, \perp) \rightarrow \{\perp, 1\}$ again determines a set of final states $F = f^{-1}[1]$ (in particular, \perp is non-final). And the morphism $i : \{\perp, *\} \rightarrow (Q, \perp)$ determines a partial initial state: either $i(*)$ lies in $Q \setminus \{\perp\}$, or no initial state is defined.
- (4) In $\mathcal{D} = \mathbf{Inv}$ let us choose $X = X_0 + \widetilde{X}_0$ and $Y = \{0, 1\}$ with $\widetilde{0} = 1$. An **Inv**-automaton is a deterministic automaton with complementary states $x \mapsto \tilde{x}$ such that (i) for every transition $p \xrightarrow{a} q$ there is a complementary transition $\tilde{p} \xrightarrow{a} \tilde{q}$ and (ii) a state q is final iff \tilde{q} is non-final.
- (5) For $\mathcal{D} = \mathbf{JSL}_0$ with $X = \mathcal{P}_f X_0$ and $Y = \{0, 1\}$ (the two-chain) an automaton consists of a semilattice Q of states, transitions $\delta_a : Q \rightarrow Q$ for $a \in X_0$ preserving finite joins (including 0), an initial state $i \in Q$ and a homomorphism $f : Q \rightarrow \{0, 1\}$ which defines a *prime upset* $F = f^{-1}[1] \subseteq Q$ of final states. Primality means that a finite join of states is final iff one of the states is. In particular, 0 is non-final. For example, the classical determinization of a nondeterministic automaton in **Set** via the powerset construction yields a **JSL**₀-automaton, where the semilattice structure is given by finite union.
- (6) More generally, automata in $\mathcal{D} = \mathbf{Mod}(\mathbb{S})$ with $X = \Psi X_0$ and $Y = \mathbb{S}$ are *\mathbb{S} -weighted automata*. Such an automaton consists of an \mathbb{S} -module Q of states, linear transitions $\delta_a : Q \rightarrow Q$ for $a \in X_0$, an initial state $i \in Q$ and a linear output map $f : Q \rightarrow \mathbb{S}$.

Remark 2.18. (1) An *algebra* for an endofunctor F of \mathcal{D} is a pair (Q, α) consisting of an object Q and a morphism $\alpha : FQ \rightarrow Q$. A *homomorphism* $h : (Q, \alpha) \rightarrow (Q', \alpha')$ of F -algebras is a morphism $h : Q \rightarrow Q'$ with $h \cdot \alpha = \alpha' \cdot Fh$. Throughout this paper we work with the endofunctor

$$FQ = I + X \otimes Q.$$

Its algebras are denoted as triples (Q, δ, i) with $\delta : X \otimes Q \rightarrow Q$ and $i : I \rightarrow Q$. Hence \mathcal{D} -automata are precisely F -algebras equipped with an output morphism $f : Q \rightarrow Y$. Moreover, automata homomorphisms are precisely F -algebra homomorphisms preserving outputs.

- (2) Analogously, a *coalgebra* for an endofunctor T of \mathcal{D} is a pair (Q, γ) consisting of an object Q and a morphism $\gamma : Q \rightarrow TQ$. A *homomorphism* $h : (Q, \gamma) \rightarrow (Q', \gamma')$ of T -coalgebras is a morphism $h : Q \rightarrow Q'$ with $Th \cdot \gamma = \gamma' \cdot h$. Throughout this paper we work with the endofunctor

$$TQ = Y \times [X, Q].$$

Its coalgebras are denoted as triples (Q, τ, f) with $\tau : Q \rightarrow [X, Q]$ and $f : Q \rightarrow Y$. Hence \mathcal{D} -automata are precisely *pointed* T -coalgebras, i.e. T -coalgebras equipped with a morphism $i : I \rightarrow Q$. Indeed, given a pointed coalgebra $I \xrightarrow{i} Q \xrightarrow{\langle f, \tau \rangle} Y \times [X, Q]$, the

morphism $Q \xrightarrow{\tau} [X, Q]$ is the curried form of a morphism $Q \otimes X \xrightarrow{\cong} X \otimes Q \xrightarrow{\delta} Q$. Hence automata homomorphisms are precisely the T -coalgebra homomorphisms preserving initial states.

Definition 2.19. Given a \mathcal{D} -monoid (M, m, i) and a morphism $e : X \rightarrow M$ of \mathcal{D} , the F -algebra associated to M and e has carrier M and structure

$$\delta \equiv (X \otimes M \xrightarrow{e \otimes M} M \otimes M \xrightarrow{m} M) \quad \text{and} \quad i : I \rightarrow M.$$

In particular, the F -algebra associated to the free monoid X^* (and its universal arrow η_X) is

$$\delta_X \equiv (X \otimes X^* \xrightarrow{\eta_X \otimes X^*} X^* \otimes X^* \xrightarrow{m_X} X^*) \quad \text{and} \quad i_X : I \rightarrow X^*.$$

Example 2.20. In **Set** every monoid M together with an ‘input’ map $e : X \rightarrow M$ determines an F -algebra with initial state i and transitions $\delta_a = e(a) \bullet -$ for all $a \in X$. The F -algebra associated to X^* is the automaton of words: its initial state is ε and the transitions are given by $w \xrightarrow{a} aw$ for $a \in X$.

Proposition 2.21 (Goguen [12]). *For any symmetric monoidal closed category \mathcal{D} with countable coproducts, X^* is the initial algebra for F .*

Remark 2.22. (1) For any F -algebra (Q, δ, i) the unique F -algebra homomorphism $e_Q : X^* \rightarrow Q$ is constructed as follows: extend the morphism $\lambda\delta : X \rightarrow [Q, Q]$ to a \mathcal{D} -monoid morphism $(\lambda\delta)^+ : X^* \rightarrow [Q, Q]$. Then

$$e_Q = (X^* \cong X^* \otimes I \xrightarrow{(\lambda\delta)^+ \otimes i} [Q, Q] \otimes Q \xrightarrow{\text{ev}} Q), \quad (2.1)$$

where **ev** is the ‘evaluation morphism’, i.e. the counit of the adjunction $- \otimes Q \dashv [Q, -]$.

To see this, recall that in any symmetric monoidal closed category \mathcal{D} the object $[Q, Q]$ can be equipped with a \mathcal{D} -monoid structure: the unit $j : I \rightarrow [Q, Q]$ and multiplication $m : [Q, Q] \otimes [Q, Q] \rightarrow [Q, Q]$ are the unique morphisms making the following diagrams commutative (where $\iota'_Q : Q \rightarrow I \otimes Q$ is the left unit isomorphism):

$$\begin{array}{ccc} [Q, Q] \otimes Q & \xrightarrow{\text{ev}} & Q \\ j \otimes Q \uparrow & \nearrow \iota'_Q & \\ I \otimes Q & & \end{array} \quad \begin{array}{ccc} [Q, Q] \otimes Q & \xrightarrow{\text{ev}} & Q \\ m \otimes Q \uparrow & & \uparrow \text{ev} \\ [Q, Q] \otimes [Q, Q] \otimes Q & \xrightarrow{[Q, Q] \otimes \text{ev}} & [Q, Q] \otimes Q \end{array}$$

Then the following commutative diagram shows that e_Q is an F -algebra homomorphism, as claimed:

$$\begin{array}{ccccc}
X \otimes X^\otimes & \xrightarrow{\eta_X \otimes X^\otimes} & X^\otimes \otimes X^\otimes & \xrightarrow{m_X} & X^\otimes \xleftarrow{i_X} I \\
\downarrow X \otimes \iota_{X^\otimes} & & \downarrow X^\otimes \otimes \iota_{X^\otimes} & & \downarrow \iota_{X^\otimes} & \downarrow \iota_I \\
X \otimes X^\otimes \otimes I & \xrightarrow{\eta_X \otimes X^\otimes \otimes I} & X^\otimes \otimes X^\otimes \otimes I & \xrightarrow{m_X \otimes I} & X^\otimes \otimes I \xleftarrow{i_X \otimes I} I \otimes I \\
\downarrow X \otimes (\lambda\delta)^+ \otimes i & & \downarrow (\lambda\delta)^+ \otimes (\lambda\delta)^+ \otimes i & & \downarrow (\lambda\delta)^+ \otimes i & \downarrow I \otimes i \\
X \otimes [Q, Q] \otimes Q & \xrightarrow{\lambda\delta \otimes [Q, Q] \otimes Q} & [Q, Q] \otimes [Q, Q] \otimes Q & \xrightarrow{m \otimes Q} & [Q, Q] \otimes Q \xleftarrow{j \otimes Q} I \otimes Q \\
\downarrow X \otimes \text{ev} & & \downarrow [Q, Q] \otimes \text{ev} & & \downarrow \text{ev} & \swarrow (\iota_Q)^{-1} \\
X \otimes Q & \xrightarrow{\lambda\delta \otimes Q} & [Q, Q] \otimes Q & \xrightarrow{\text{ev}} & Q & \xleftarrow{i} I \otimes Q
\end{array}$$

δ

- (2) If \mathcal{D} is a commutative variety of algebras or ordered algebras, the monoid structure on $[Q, Q]$ is given by composition and the identity map. Let $\delta_x : Q \rightarrow Q$ denote the image of $x \in X^\otimes$ under $(\lambda\delta)^+ : X^\otimes \rightarrow [Q, Q]$. Then the initial F -algebra homomorphism e_Q in (2.1) sends x to the state $\delta_x \cdot i : I \rightarrow Q$. This follows from the commutative diagram below, see Remark 2.6:

$$\begin{array}{ccccc}
|X^\otimes| & \xrightarrow{\cong} & |X^\otimes \otimes I| & \xrightarrow{(\lambda\delta)^+ \otimes i} & |[Q, Q] \otimes Q| & \xrightarrow{\text{ev}} & |Q| \\
\searrow \langle X^\otimes, c_1 \rangle & & \uparrow \eta & & \uparrow \eta & \swarrow \text{ev}' & \\
& & |X^\otimes| \times |I| & \xrightarrow{(\lambda\delta)^+ \times i} & |[Q, Q]| \times |Q| & &
\end{array}$$

e_Q

Notation 2.23. $\delta^\otimes : X^\otimes \otimes Q \rightarrow Q$ denotes the uncurried form of $(\lambda\delta)^+ : X^\otimes \rightarrow [Q, Q]$.

Remark 2.24. (1) One can also define δ^\otimes explicitly (using that $- \otimes Q$ preserves the coproduct $X^\otimes = \coprod_{n < \omega} X^{\otimes n}$) as $\delta^\otimes = [\delta_n^\otimes]_{n < \omega}$ where $\delta_0^\otimes = (I \otimes Q \xrightarrow{\cong} Q)$ and

$$\delta_{n+1}^\otimes = (X^{\otimes(n+1)} \otimes Q = X \otimes X^{\otimes n} \otimes Q \xrightarrow{X \otimes \delta_n^\otimes} X \otimes Q \xrightarrow{\delta} Q).$$

From this definition we clearly have the equation

$$\delta = (X \otimes Q \xrightarrow{\eta_X \otimes Q} X^\otimes \otimes Q \xrightarrow{\delta^\otimes} Q).$$

- (2) By [12, Theorem 3.3] the morphism $\delta^\otimes : X^\otimes \otimes Q \rightarrow Q$ defines an *action* of the monoid X^\otimes on Q , i.e. the following diagrams commute:

$$\begin{array}{ccc}
I \otimes Q & \xrightarrow{i_X \otimes Q} & X^\otimes \otimes Q \\
\cong \searrow & & \downarrow \delta^\otimes \\
& & Q
\end{array}
\quad \text{and} \quad
\begin{array}{ccc}
X^\otimes \otimes X^\otimes \otimes Q & \xrightarrow{m_X \otimes Q} & X^\otimes \otimes Q \\
X^\otimes \otimes \delta^\otimes \downarrow & & \downarrow \delta^\otimes \\
X^\otimes \otimes Q & \xrightarrow{\delta^\otimes} & Q
\end{array}$$

Moreover, every F -algebra homomorphism h from (Q, δ, i) to (Q', δ', i') is a morphism of the corresponding monoid actions, i.e. the following diagram commutes:

$$\begin{array}{ccc} X^\otimes \otimes Q & \xrightarrow{\delta^\otimes} & Q \xleftarrow{i} I \\ X^\otimes \otimes h \downarrow & & \downarrow h \swarrow i' \\ X^\otimes \otimes Q' & \xrightarrow{(\delta')^\otimes} & Q' \end{array}$$

- (3) Furthermore, it is not difficult to prove that for $\delta_X : X \otimes X^\otimes \rightarrow X^\otimes$ coming from the F -algebra associated to X^\otimes we have that δ_X^\otimes is the monoid multiplication:

$$\delta_X^\otimes = m_X : X^\otimes \otimes X^\otimes \rightarrow X^\otimes.$$

Remark 2.25. Recall from Rutten [21] that the final coalgebra for the functor $TQ = \{0, 1\} \times Q^X$ on **Set** is carried by the set $\mathcal{P}X^* \cong [X^*, \{0, 1\}]$ of all languages over X . The transitions are given by taking *left derivatives*,

$$L \xrightarrow{a} a^{-1}L = \{w \in X^* : aw \in L\} \quad \text{for } L \in \mathcal{P}X^* \text{ and } a \in X,$$

and the output map $\mathcal{P}X^* \rightarrow \{0, 1\}$ sends L to 1 iff L contains the empty word. Given any coalgebra Q , the unique coalgebra homomorphism from Q to $\mathcal{P}\Sigma^*$ assigns to every state q the language accepted by q (as an initial state).

These observations generalize to arbitrary symmetric monoidal closed categories. The object $[X^\otimes, Y]$ of \mathcal{D} carries the following T -coalgebra structure: its transition morphism $\tau_{[X^\otimes, Y]} : [X^\otimes, Y] \rightarrow [X, [X^\otimes, Y]]$ is the two-fold currying of

$$[X^\otimes, Y] \otimes X \otimes X^\otimes \xrightarrow{[X^\otimes, Y] \otimes \eta_X \otimes X^\otimes} [X^\otimes, Y] \otimes X^\otimes \otimes X^\otimes \xrightarrow{[X^\otimes, Y] \otimes m_X} [X^\otimes, Y] \otimes X^\otimes \xrightarrow{\text{ev}} Y,$$

and its output morphism $f_{[X^\otimes, Y]} : [X^\otimes, Y] \rightarrow Y$ is

$$f_{[X^\otimes, Y]} = ([X^\otimes, Y] \cong [X^\otimes, Y] \otimes I \xrightarrow{[X^\otimes, Y] \otimes i_X} [X^\otimes, Y] \otimes X^\otimes \xrightarrow{\text{ev}} Y).$$

Proposition 2.26. *For any symmetric monoidal closed category \mathcal{D} , the coalgebra $[X^\otimes, Y]$ is the final coalgebra for T .*

Proof. Given a T -coalgebra (Q, τ, f) , consider the morphism $\delta = (X \otimes Q \xrightarrow{\cong} Q \otimes X \xrightarrow{\beta} Q)$ where β is the uncurried form of $\tau : Q \rightarrow [X, Q]$, and denote by $\delta^\otimes : X^\otimes \otimes Q \rightarrow Q$ its extension, see Notation 2.23. We claim that the unique coalgebra homomorphism into $[X^\otimes, Y]$ is $\lambda h : Q \rightarrow [X^\otimes, Y]$, where

$$h = (Q \otimes X^\otimes \cong X^\otimes \otimes Q \xrightarrow{\delta^\otimes} Q \xrightarrow{f} Y). \quad (2.2)$$

Let us first prove that h is indeed a coalgebra homomorphism. Preservation of outputs is shown by the following commutative diagram (for the upper left-hand part use Remark 2.24(2)):

$$\begin{array}{ccccc}
 Q & \xrightarrow{\quad\quad\quad} & Q & \xrightarrow{f} & Y \\
 \parallel & \searrow & \uparrow \delta^\otimes & \nearrow h & \uparrow \text{ev} \\
 Q & \xrightarrow{\cong} & I \otimes Q & \xrightarrow{i_X \otimes Q} & X^\otimes \otimes Q \\
 \parallel & \searrow & \uparrow \cong & \nearrow & \\
 Q & \xrightarrow{\cong} & Q \otimes I & \xrightarrow{Q \otimes i_X} & Q \otimes X^\otimes \\
 \downarrow \lambda h & & \downarrow \lambda h \otimes I & \searrow \lambda h \otimes X^\otimes & \\
 [X^\otimes, Y] & \xrightarrow{\cong} & [X^\otimes, Y] \otimes I & \xrightarrow{[X^\otimes, Y] \otimes i_X} & [X^\otimes, Y] \otimes X^\otimes
 \end{array}$$

For preservation of transitions it suffices to show that the following diagram commutes, where $\bar{\tau} : [X^\otimes, Y] \otimes X \rightarrow [X^\otimes, Y]$ is the uncurried coalgebra structure of $[X^\otimes, Y]$:

$$\begin{array}{ccc}
 Q \otimes X & \xrightarrow{\beta} & Q \\
 \lambda h \otimes X \downarrow & & \downarrow \lambda h \\
 [X^\otimes, Y] \otimes X & \xrightarrow{\bar{\tau}} & [X^\otimes, Y]
 \end{array}$$

But the above diagram is precisely the curried version of the following one, which commutes by the properties listed in Remark 2.24. (We omit writing \otimes for space reasons.)

$$\begin{array}{ccccccc}
 QX^\otimes & \xrightarrow{\beta X^\otimes} & & & & & QX^\otimes \\
 \parallel & \searrow & \nearrow \delta X^\otimes & \nearrow \delta^\otimes X^\otimes & \nearrow \cong & \searrow & \downarrow h \\
 QX^\otimes & \xrightarrow{Q\eta_X X^\otimes} & XQX^\otimes & \xrightarrow{\eta_X QX^\otimes} & X^\otimes QX^\otimes & \xrightarrow{\cong} & X^\otimes X^\otimes Q & \xrightarrow{X^\otimes \delta^\otimes} & X^\otimes Q \\
 \downarrow \lambda h X X^\otimes & \searrow & \downarrow Q\eta_X X^\otimes & \nearrow \cong & \downarrow \delta^\otimes & \searrow & \downarrow \delta^\otimes & \searrow & \\
 QX^\otimes & \xrightarrow{Q\eta_X X^\otimes} & QX^\otimes X^\otimes & \xrightarrow{Qm_X} & QX^\otimes & \xrightarrow{\cong} & X^\otimes Q & \xrightarrow{\delta^\otimes} & Q \\
 \downarrow [X^\otimes, Y] \eta_X X^\otimes & \searrow & \downarrow \lambda h X^\otimes X^\otimes & \nearrow [X^\otimes, Y] m_X & \downarrow \lambda h X^\otimes & \searrow & \downarrow h & \searrow & \\
 [X^\otimes, Y] X X^\otimes & \xrightarrow{[X^\otimes, Y] \eta_X X^\otimes} & [X^\otimes, Y] X^\otimes X^\otimes & \xrightarrow{[X^\otimes, Y] m_X} & [X^\otimes, Y] X^\otimes & \xrightarrow{\text{ev}} & Y & &
 \end{array}$$

For the uniqueness, suppose that any coalgebra homomorphism $\lambda h : Q \rightarrow [X^\otimes, Y]$ is given. We show that $h : Q \otimes X^\otimes \rightarrow Y$ is determined by the composites $(Q \otimes X^{\otimes n} \xrightarrow{Q \otimes i_n} Q \otimes X^\otimes \xrightarrow{h} Y)$, $n < \omega$, where $i_n : X^{\otimes n} \rightarrow X^\otimes$ is the n -th coproduct injection. This proves the uniqueness of λh : since \otimes preserves coproducts, the morphisms $(Q \otimes i_n)_{n < \omega}$ form a coproduct cocone.

For $n = 0$, the claim is proved by the following diagram:

$$\begin{array}{ccc}
 Q \otimes I & \xrightarrow{\cong} & Q \\
 \downarrow Q \otimes i_0 & \searrow \lambda h \otimes I & \swarrow \lambda h \\
 & [X^\otimes, Y] \otimes I & \xrightarrow[\cong]{\cong} [X^\otimes, Y] \\
 & \downarrow [X^\otimes, Y] \otimes i_0 & \downarrow f \\
 & [X^\otimes, Y] \otimes X^\otimes & \xrightarrow{f_{[X^\otimes, Y]}} Y \\
 \downarrow \lambda h \otimes X^\otimes & \nearrow & \downarrow \\
 Q \otimes X^\otimes & \xrightarrow{h} & Y
 \end{array}$$

And the following diagram shows that $h \cdot (Q \otimes i_{n+1})$ is determined by $h \cdot (Q \otimes i_n)$ (again we omit \otimes , in particular we write X^n for $X^{\otimes n}$):

$$\begin{array}{ccccc}
 QXX^n & \xrightarrow{Q i_{n+1}} & QX^\otimes & \xrightarrow{h} & Y \\
 \downarrow \beta X^n & \searrow \lambda h XX^n & \downarrow \lambda h X^\otimes & \nearrow \text{ev} & \downarrow h \\
 & [X^\otimes, Y]XX^n & \xrightarrow{[X^\otimes, Y] i_{n+1}} & [X^\otimes, Y]X^\otimes & \\
 & \downarrow \bar{\tau} X^n & \searrow [X^\otimes, Y] X i_n & \nearrow [X^\otimes, Y] \eta_X X^\otimes & \\
 & [X^\otimes, Y]XX^\otimes & \xrightarrow{[X^\otimes, Y] X^\otimes X^\otimes} & [X^\otimes, Y]X^\otimes X^\otimes & \\
 & \downarrow \bar{\tau} X^\otimes & \searrow & \nearrow (*) & \\
 & [X^\otimes, Y]X^n & \xrightarrow{[X^\otimes, Y] i_n} & [X^\otimes, Y]X^\otimes & \\
 \downarrow \lambda h X^n & \nearrow & \downarrow \lambda h X^\otimes & \nearrow \text{ev} & \\
 QX^n & \xrightarrow{Q i_n} & QX^\otimes & & Y
 \end{array}$$

Note that part marked with $(*)$ commutes by the definition of the coalgebra structure on $[X^\otimes, Y]$ and all other parts are easy to see. \square

Remark 2.27. If \mathcal{D} is a commutative variety of algebras or ordered algebras, the final homomorphism $\lambda h : Q \rightarrow [X^\otimes, Y]$ sends a state $q \in |Q|$ to the morphism $x \mapsto f \cdot \delta_x(q)$, with $\delta_x : Q \rightarrow Q$ defined as in Remark 2.22(2). To see this, consider the commutative diagram below, where $(\delta^\otimes)'$ is the bimorphism corresponding to δ^\otimes :

$$\begin{array}{ccccc}
 & & \xrightarrow{h} & & \\
 |Q \otimes X^\otimes| & \xrightarrow{\cong} & |X^\otimes \otimes Q| & \xrightarrow{\delta^\otimes} & |Q| \xrightarrow{f} |Y| \\
 \eta \uparrow & & \eta \uparrow & \nearrow (\delta^\otimes)' & \\
 |Q| \times |X^\otimes| & \xrightarrow{\cong} & |X^\otimes| \times |Q| & &
 \end{array}$$

By the definition of δ_x and Remark 2.6, the bimorphism $(\delta^\otimes)'$ maps (x, q) to $\delta_x(q)$. Thus $h \cdot \eta$ maps (q, x) to $f \cdot \delta_x(q)$. This implies the claim since λh is the curried form of $h \cdot \eta$.

Proposition 2.26 motivates the following definition:

Definition 2.28 (Goguen [12]). A *language* in \mathcal{D} is a morphism $L : X^\otimes \rightarrow Y$.

Note that if $X = \Psi X_0$ (and hence $X^\otimes = \Psi X_0^*$) for some set X_0 , one can identify a language $L : X^\otimes \rightarrow Y$ in \mathcal{D} with its adjoint transpose $\tilde{L} : X_0^* \rightarrow |Y|$, using the adjunction $\Psi \dashv |-| : \mathcal{D} \rightarrow \mathbf{Set}$. In the case where $|Y|$ is a two-element set, \tilde{L} is the characteristic function of a “classical” language $L_0 \subseteq X_0^*$.

- Example 2.29.** (1) In $\mathcal{D} = \mathbf{Set}$ (with $X^\otimes = X^*$ and $Y = \{0, 1\}$) one represents $L_0 \subseteq X^*$ by its characteristic function $L : X^* \rightarrow \{0, 1\}$. Analogously for $\mathcal{D} = \mathbf{Pos}$ with $Y = \{0 < 1\}$.
(2) In $\mathcal{D} = \mathbf{Set}_\perp$ (with $X = X_0 + \{\perp\}$, $X^\otimes = X_0^* + \{\perp\}$ and $Y = \{\perp, 1\}$) one represents $L_0 \subseteq X_0^*$ by its extended characteristic function $L : X_0^* + \{\perp\} \rightarrow \{\perp, 1\}$ where $L(\perp) = \perp$.
(3) In $\mathcal{D} = \mathbf{Inv}$ (with $X = X_0 + \widetilde{X}_0$, $X^\otimes = X_0^* + \widetilde{X}_0^*$ and $Y = \{0, 1\}$) one represents $L_0 \subseteq X_0^*$ by $L : X_0^* + \widetilde{X}_0^* \rightarrow \{0, 1\}$ where $L(w) = 1$ iff $w \in L_0$ and $L(\tilde{w}) = 1$ iff $w \notin L_0$ for all words $w \in X_0^*$.
(4) In $\mathcal{D} = \mathbf{JSL}_0$ (with $X = \mathcal{P}_f X_0$, $X^\otimes = \mathcal{P}_f X_0^*$ and $Y = \{0, 1\}$) one represents $L_0 \subseteq X_0^*$ by $L : \mathcal{P}_f X_0^* \rightarrow \{0, 1\}$ where $L(U) = 1$ iff $U \cap L_0 \neq \emptyset$.
(5) In $\mathcal{D} = \mathbf{Mod}(\mathbb{S})$ (with $X = \Psi X_0$, $X^\otimes = \mathbb{S}[X_0]$ and $Y = \mathbb{S}$) an \mathbb{S} -weighted language $L_0 : X_0^* \rightarrow \mathbb{S}$ is represented by its free extension to a module homomorphism

$$L : \mathbb{S}[X_0] \rightarrow \mathbb{S}, \quad L \left(\sum_{i=1}^n c(w_i) w_i \right) = \sum_{i=1}^n c(w_i) L_0(w_i).$$

Definition 2.30 (Goguen [12]). The language *accepted* by a \mathcal{D} -automaton (Q, δ, i, f) is $L_Q = (X^\otimes \xrightarrow{e_Q} Q \xrightarrow{f} Y)$, where e_Q is the F -algebra homomorphism of Remark 2.22.

- Example 2.31.** (1) In $\mathcal{D} = \mathbf{Set}$ or \mathbf{Pos} with $Y = \{0, 1\}$, the homomorphism $e_Q : X^* \rightarrow Q$ assigns to every word w the state it computes in Q , i.e. the state the automaton reaches on input w . Thus $L_Q(w) = 1$ iff Q terminates in a final state on input w , which is precisely the standard definition of the accepted language of an (ordered) automaton. For general Y , the function $L_Q : X^* \rightarrow Y$ is the behavior of the (ordered) Moore automaton Q , i.e. $L_Q(w)$ is the output of the last state in the computation of w .
(2) For $\mathcal{D} = \mathbf{Set}_\perp$ with $X = X_0 + \{\perp\}$ and $Y = \{\perp, 1\}$, we have $e_Q : X_0^* + \{\perp\} \rightarrow (Q, \perp)$ sending \perp to \perp , and sending a word in X_0^* to the state it computes (if any), and to \perp otherwise. Hence $L_Q : X_0^* + \{\perp\} \rightarrow \{\perp, 1\}$ defines (via the preimage of 1) the usual language accepted by a partial automaton.
(3) In $\mathcal{D} = \mathbf{Inv}$ with $X = X_0 + \widetilde{X}_0$ and $Y = \{0, 1\}$, the map $L_Q : X_0^* + \widetilde{X}_0^* \rightarrow \{0, 1\}$ sends $w \in X_0^*$ to 1 iff w computes a final state, and it sends $\tilde{w} \in \widetilde{X}_0^*$ to 1 iff w computes a non-final state.
(4) In $\mathcal{D} = \mathbf{JSL}_0$ with $X = \mathcal{P}_f X_0$ and $Y = \{0, 1\}$, the map $L_Q : \mathcal{P}_f X_0^* \rightarrow \{0, 1\}$ assigns to $U \in \mathcal{P}_f X_0^*$ the value 1 iff the computation of at least one word in U ends in a final state.
(5) In $\mathcal{D} = \mathbf{Mod}(\mathbb{S})$ with $X = \Psi X_0$ and $Y = \mathbb{S}$, the map $L_Q : \mathbb{S}[X_0] \rightarrow \mathbb{S}$ assigns to $\sum_{i=1}^n c(w_i) w_i$ the value $\sum_{i=1}^n c(w_i) y_i$, where y_i is the output of the state the automaton computes on input w_i . Taking $Q = \mathbb{S}^n$ for some natural number n yields a classical n -state weighted automaton; indeed, $i \in \mathbb{S}^n$ is an initial vector, the linear map $f : \mathbb{S}^n \rightarrow \mathbb{S}$ corresponds to an output vector $o \in \mathbb{S}^n$ and the linear transition map $\delta : X \otimes \mathbb{S}^n \rightarrow \mathbb{S}^n$ is given by a family of linear maps $(\delta_a : \mathbb{S}^n \rightarrow \mathbb{S}^n)_{a \in X_0}$, which can be represented by a family $(M_a)_{a \in X_0}$ of $n \times n$ matrices over \mathbb{S} . It then follows that the restriction of L_Q to a map $X_0^* \rightarrow \mathbb{S}$ is the usual weighted language assigning to a word w the element $o \cdot M_w \cdot i^T$ of \mathbb{S} , where M_w is the obvious product of the matrices M_a .

Remark 2.32. By Remark 2.18 every \mathcal{D} -automaton (Q, δ, i, f) defines an F -algebra as well as a T -coalgebra. Our above definition of L_Q was purely algebraic. The corresponding coalgebraic definition uses the unique coalgebra homomorphism $c_Q : Q \rightarrow [X^\otimes, Y]$ into the final T -coalgebra and precomposes with $i : I \rightarrow Q$ to get a morphism $c_Q \cdot i : I \rightarrow [X^\otimes, Y]$ (choosing a language, i.e. an element of $[X^\otimes, Y]$). Unsurprisingly, the results are equal:

Proposition 2.33. *The language $L_Q : X^\otimes \rightarrow Y$ of an automaton (Q, δ, i, f) is the uncurried form of the morphism $c_Q \cdot i : I \rightarrow [X^\otimes, Y]$.*

Proof. Recall from the proof of Proposition 2.26 that the uncurried version of c_Q is the morphism

$$Q \otimes X^\otimes \cong X^\otimes \otimes Q \xrightarrow{\delta^\otimes} Q \xrightarrow{f} Y.$$

Hence $c_Q \cdot i : I \rightarrow [X^\otimes, Y]$ determines the language

$$X^\otimes \cong X^\otimes \otimes I \xrightarrow{X^\otimes \otimes i} X^\otimes \otimes Q \xrightarrow{\delta^\otimes} Q \xrightarrow{f} Y,$$

and this is precisely L_Q , as shown by the diagram below (where δ_X is given by the F -algebra structure associated to X^\otimes and η_X , see Remark 2.19):

$$\begin{array}{ccccccc}
& & & \text{id} & & & \\
& & & \curvearrowright & & & \\
X^\otimes & \xrightarrow{\cong} & X^\otimes \otimes I & \xrightarrow{X^\otimes \otimes i_X} & X^\otimes \otimes X^\otimes & \xrightarrow{\delta_X^\otimes} & X^\otimes & \xrightarrow{L_Q} & Y \\
& & \searrow^{X^\otimes \otimes i} & & \downarrow^{X^\otimes \otimes e_Q} & & \downarrow^{e_Q} & & \\
& & & & X^\otimes \otimes Q & \xrightarrow{\delta^\otimes} & Q & \xrightarrow{f} & Y
\end{array}$$

Indeed, the right-hand triangle commutes by the definition of L_Q and the left-hand one and the inner square commute since e_Q is an F -algebra homomorphism (see Remark 2.24(2)). The upper part commutes since δ_X^\otimes is the monoid multiplication m_X , see Remark 2.24(3). \square

3. ALGEBRAIC RECOGNITION AND SYNTACTIC \mathcal{D} -MONOIDS

In classical algebraic automata theory one considers recognition of languages by (ordinary) monoids in lieu of automata. The key concept is the *syntactic monoid* which is characterized as the smallest monoid recognizing a given language. There are also related concepts of canonical algebraic recognizers in the literature, e.g. the syntactic ordered monoid, the syntactic idempotent semiring and the syntactic associative algebra. In this section we will give a uniform account of algebraic language recognition in our categorical setting. Our main result is the definition and construction of a minimal algebraic recognizer, the *syntactic \mathcal{D} -monoid* of a language.

Assumptions 3.1. Throughout this section $(\mathcal{D}, \otimes, I)$ is an arbitrary symmetric monoidal closed category, and \mathcal{E} is a class of epimorphisms in \mathcal{D} that contains all isomorphisms and is closed under composition. In the case where \mathcal{D} is a variety of algebras or ordered algebras, we always choose

$$\mathcal{E} = \text{surjective homomorphisms.}$$

Definition 3.2. A \mathcal{D} -monoid morphism $e : X^\otimes \rightarrow M$ recognizes the language $L : X^\otimes \rightarrow Y$ if there exists a morphism $f : M \rightarrow Y$ of \mathcal{D} with $L = f \cdot e$.

Example 3.3. We use the notation of Example 2.29.

- (1) $\mathcal{D} = \mathbf{Set}$ with $X^\otimes = X^*$ and $Y = \{0, 1\}$: given a monoid M , a function $f : M \rightarrow \{0, 1\}$ defines a subset $F = f^{-1}[1] \subseteq M$. Hence a monoid morphism $e : X^* \rightarrow M$ recognizes L via f (i.e. $L = f \cdot e$) iff $L_0 = e^{-1}[F]$. This is the classical notion of recognition of a language $L_0 \subseteq X^*$ by a monoid, see e.g. [15].
- (2) $\mathcal{D} = \mathbf{Pos}$ with $X^\otimes = X^*$ and $Y = \{0 < 1\}$: given an ordered monoid M , a monotone map $f : M \rightarrow \{0, 1\}$ defines an upper set $F = f^{-1}[1] \subseteq M$. Hence a monoid morphism $e : X^* \rightarrow M$ recognizes L iff L_0 is the preimage of some upper set of M . This notion of recognition is due to Pin [16].
- (3) $\mathcal{D} = \mathbf{Set}_\perp$ with $X = X_0 + \{\perp\}$, $X^\otimes = X_0^* + \{\perp\}$ and $Y = \{\perp, 1\}$: given a monoid with zero M , a \mathbf{Set}_\perp -morphism $f : M \rightarrow \{\perp, 1\}$ defines a subset $F = f^{-1}[1]$ of $M \setminus \{0\}$. A zero-preserving monoid morphism $e : X_0^* + \{\perp\} \rightarrow M$ recognizes L via f iff $L_0 = e^{-1}[F]$.
- (4) $\mathcal{D} = \mathbf{Inv}$ with $X = X_0 + \widetilde{X}_0$, $X^\otimes = X_0^* + \widetilde{X}_0^*$ and $Y = \{0, 1\}$: for an involution monoid M to give a morphism $f : M \rightarrow \{0, 1\}$ means to give a subset $F = f^{-1}[1] \subseteq M$ satisfying $m \in F$ iff $\widetilde{m} \notin F$. Then L is recognized by $e : X_0^* + \widetilde{X}_0^* \rightarrow M$ via f iff $L_0 = X_0^* \cap e^{-1}[F]$.
- (5) $\mathcal{D} = \mathbf{JSL}_0$ with $X = \mathcal{P}_f X_0$, $X^\otimes = \mathcal{P}_f X_0^*$ and $Y = \{0, 1\}$: for an idempotent semiring M a morphism $f : M \rightarrow Y$ defines a prime upset $F = f^{-1}[1]$, see Example 2.17. Hence L is recognized by a semiring homomorphism $e : \mathcal{P}_f X_0^* \rightarrow M$ via f iff $L_0 = X_0^* \cap e^{-1}[F]$. Here we identify X_0^* with the set of all singleton languages $\{w\}$, $w \in X_0^*$. This is the concept of language recognition introduced by Polák [18] (except that he puts $F = f^{-1}[0]$, so 0 and 1 must be swapped, as well as F and $M \setminus F$).
- (6) $\mathcal{D} = \mathbf{Mod}(\mathbb{S})$ with $X = \Psi X_0$, $X^\otimes = \mathbb{S}[X_0]$ and $Y = \mathbb{S}$: given an associative algebra M , a language L is recognized by $e : \mathbb{S}[X_0] \rightarrow M$ via $f : M \rightarrow \mathbb{S}$ iff $L = f \cdot e$. For the case where the semiring \mathbb{S} is a ring, this notion of recognition is due to Reutenauer [20].

Remark 3.4. (1) By an X -generated \mathcal{D} -monoid we mean a morphism $e : X^\otimes \twoheadrightarrow M$ in $\mathbf{Mon}(\mathcal{D})$ with $e \in \mathcal{E}$. Given two X -generated \mathcal{D} -monoids $e_i : X^\otimes \twoheadrightarrow M_i$, $i = 1, 2$, we say, as usual, that e_1 is *smaller or equal to* e_2 (notation: $e_1 \leq e_2$) if e_1 factorizes through e_2 . Note that if \mathcal{D} is a variety and $X = \Psi X_0$, the free \mathcal{D} -monoid $X^\otimes = \Psi X_0^*$ on X is also the free \mathcal{D} -monoid on the set X_0 (w.r.t. the forgetful functor $\mathbf{Mon}(\mathcal{D}) \rightarrow \mathbf{Set}$), see Proposition 2.14. In this case, to give a quotient $e : X^\otimes \twoheadrightarrow M$ is equivalent to giving an X_0 -indexed family $(m_x)_{x \in X_0}$ of generators for the \mathcal{D} -monoid M – which is why M may also be called an X_0 -generated \mathcal{D} -monoid.

- (2) Let $e : X^\otimes \twoheadrightarrow M$ be an X -generated \mathcal{D} -monoid with unit $i : I \rightarrow M$ and multiplication $m : M \otimes M \rightarrow M$. Recall that $\eta_X : X \rightarrow X^\otimes$ denotes the universal morphism of the free \mathcal{D} -monoid on X and consider the F -algebra associated to M and $X \xrightarrow{\eta_X} X^\otimes \xrightarrow{e} M$ (see Definition 2.19). Thus, together with a given $f : M \rightarrow Y$ an X -generated \mathcal{D} -monoid induces an automaton (M, δ, i, f) called the *derived automaton*.

Lemma 3.5. *The language recognized by an X -generated \mathcal{D} -monoid $e : X^\otimes \twoheadrightarrow M$ via $f : M \rightarrow Y$ is the language accepted by its derived automaton.*

Proof. By definition e recognizes via f the language $L = f \cdot e$. We are done once we prove that e is the unique F -algebra homomorphism from X^\otimes to the F -algebra associated to M and $e \cdot \eta_X$ (cf. Remark 3.4). Recall from Proposition 2.21 that the initial F -algebra is the F -algebra associated to the free \mathcal{D} -monoid X^\otimes and η_X . Then the following diagram clearly

commutes, since e is a \mathcal{D} -monoid morphism:

$$\begin{array}{ccccc}
 FX^{\otimes} = I + X \otimes X^{\otimes} & \xrightarrow{I + \eta_X \otimes X^{\otimes}} & I + X^{\otimes} \otimes X^{\otimes} & \xrightarrow{[i_X, m_X]} & X^{\otimes} \\
 \downarrow Fe = I + X \otimes e & & \downarrow I + e \otimes e & & \downarrow e \\
 FM = I + X \otimes M & \xrightarrow{I + (e \cdot \eta_X) \otimes M} & I + M \otimes M & \xrightarrow{[i, m]} & M
 \end{array}$$

This completes the proof. \square

We are now ready to give an abstract account of syntactic monoids in our setting. In classical algebraic automata theory the syntactic monoid of a language is characterized as the smallest monoid recognizing that language. We will use this property as our definition of the syntactic \mathcal{D} -monoid.

Definition 3.6. The *syntactic \mathcal{D} -monoid* of a language $L : X^{\otimes} \rightarrow Y$, denoted by $\text{Syn}(L)$, is the smallest X -generated monoid recognizing L .

In more detail, the syntactic \mathcal{D} -monoid of L is an X -generated \mathcal{D} -monoid $e_L : X^{\otimes} \rightarrow \text{Syn}(L)$ together with a morphism $f_L : \text{Syn}(L) \rightarrow Y$ of \mathcal{D} such that (i) e_L recognizes L via f_L , and (ii) for every X -generated \mathcal{D} -monoid $e : X^{\otimes} \rightarrow M$ recognizing L via $f : M \rightarrow Y$ we have $e_L \leq e$, that is, the left-hand triangle below commutes for some \mathcal{D} -monoid morphism h :

$$\begin{array}{ccccc}
 X^{\otimes} & \xrightarrow{e} & M & \xrightarrow{f} & Y \\
 & \searrow e_L & \downarrow h & \nearrow f_L & \\
 & & \text{Syn}(L) & &
 \end{array}$$

Note that the right-hand triangle also commutes since e is epimorphic and $f \cdot e = L = f_L \cdot e_L$. The universal property determines $\text{Syn}(L)$, e_L and f_L uniquely up to isomorphism.

The above definition leaves open the question whether the syntactic \mathcal{D} -monoid of a language actually exists. In Section 3.1 we investigate the existence of syntactic \mathcal{D} -monoids in an abstract symmetric monoidal closed category \mathcal{D} . In Section 3.2 we show how to construct them in our algebraic setting, using the syntactic congruence of a language.

3.1. Existence of syntactic \mathcal{D} -monoids.

Definition 3.7. The category \mathcal{D} is *stable* w.r.t. \mathcal{E} if the following square is a pushout for all morphisms $a : A \rightarrow A'$ and $b : B \rightarrow B'$ in \mathcal{E} :

$$\begin{array}{ccccc}
 & & A \otimes B & & \\
 & \swarrow a \otimes B & \downarrow a \otimes b & \searrow A \otimes b & \\
 A' \otimes B & & & & A \otimes B' \\
 & \searrow A' \otimes b & \downarrow & \swarrow a \otimes B & \\
 & & A' \otimes B' & &
 \end{array}$$

Example 3.8. (1) **Set** is stable w.r.t. $\mathcal{E} =$ surjective maps. The pushout of the surjections $a \times B$ and $A \times b$ is given by $A \times B / \sim$ where \sim is the least equivalence relation such that

$$a(x) = a(x') \text{ implies } (x, y) \sim (x', y) \text{ for all } y \in B$$

as well as

$$b(y) = b(y') \text{ implies } (x, y) \sim (x, y') \text{ for all } x \in A.$$

Obviously, \sim is the kernel equivalence of $a \times b$.

- (2) More generally, every variety \mathcal{D} of algebras is stable w.r.t. $\mathcal{E} =$ surjective homomorphisms. To see this, recall from Definition 2.4 that the forgetful functor from \mathcal{D} to \mathbf{Set} yields the tensor product $A \otimes B$ via the universal bimorphism $\eta_{A,B} : |A| \times |B| \rightarrow |A \otimes B|$. Suppose that $u : A' \otimes B \rightarrow C$ and $v : A \otimes B' \rightarrow C$ are given with $u \cdot (a \otimes B) = v \cdot (A \otimes b)$. Consider the following diagram:

$$\begin{array}{ccccc}
 & & |A \otimes B| & & \\
 & & \uparrow \eta & & \\
 & & |A| \times |B| & & \\
 & \swarrow a \otimes B & & \searrow A \otimes b & \\
 |A' \otimes B| & \xleftarrow{\eta} & |A'| \times |B| & & |A \otimes B'| \\
 & \swarrow a \times B & \downarrow a \times b & \searrow A \times b & \\
 & & |A'| \times |B'| & & \\
 & \swarrow A' \times b & \downarrow \eta & \searrow a \times B' & \\
 & & |A' \otimes B'| & & \\
 & \swarrow A' \otimes b & \downarrow w & \searrow a \otimes B' & \\
 & & |C| & & \\
 & \swarrow u & & \searrow v &
 \end{array}$$

By Example 3.8(1) above the inner square is a pushout in \mathbf{Set} , so there is a unique function $w_0 : |A'| \times |B'| \rightarrow |C|$ with $u \cdot \eta_{A',B} = w_0 \cdot (A' \times b)$ and $v \cdot \eta_{A,B'} = w_0 \cdot (a \times B')$. We claim that w_0 is a bimorphism of \mathcal{D} . Indeed, for each $x \in |A'|$ the map $w_0(x, -)$ carries a morphism of \mathcal{D} : choosing $\bar{x} \in |A|$ with $a(\bar{x}) = x$, we have

$$w_0(x, -) = w_0 \cdot (a \times B')(\bar{x}, -) = v \cdot \eta_{A,B'}(\bar{x}, -)$$

and the last map is a morphism in \mathcal{D} because $\eta_{A,B'}$ is a bimorphism. Symmetrically, $w_0(-, y)$ is a morphism of \mathcal{D} for all $y \in |B'|$.

Consequently, the bimorphism w_0 induces a unique morphism $w : A' \otimes B' \rightarrow C$ in \mathcal{D} with $w \cdot \eta_{A',B'} = w_0$. We have

$$w \cdot (A' \otimes b) = u$$

because, by definition of w_0 , this holds when precomposed with $\eta_{A',B}$. Analogously,

$$w \cdot (a \otimes B') = v.$$

The uniqueness of w follows from the uniqueness of w_0 .

- (3) \mathbf{Pos} is stable w.r.t. $\mathcal{E} =$ surjective monotone maps. The pushout of $a \times B$ and $A \times b$ is obtained from the pushout in \mathbf{Set} , i.e. $|A'| \times |B'|$, by taking the smallest preorder \leq such that $(x, y) \leq (x', y)$ for $x \leq x'$ in A' as well as $(x, y) \leq (x, y')$ for all $y \leq y'$ in B' (and forming the corresponding quotient poset of the preordered set $A' \times B'$, cf. Remark 3.16 below). But this is just the product order; thus the pushout is the product $A' \times B'$ in \mathbf{Pos} with the cocone $A' \times b$ and $a \times B'$.

- (4) More generally, every variety \mathcal{D} of ordered algebras is stable w.r.t. $\mathcal{E} =$ surjective homomorphisms. The argument is completely analogous to Example 3.8(2).

Theorem 3.9. *Let \mathcal{D} be stable w.r.t. \mathcal{E} , and suppose moreover that \mathcal{E} is closed under wide pushouts and tensor products. Then the forgetful functor from $\mathbf{Mon}(\mathcal{D})$ to \mathcal{D} creates wide pushouts of \mathcal{E} -carried morphisms.*

Proof. (1) Given $e_i : D \rightarrow D_i$ ($i \in I$) in \mathcal{E} with wide pushout $e = \bar{e}_i \cdot e_i : D \rightarrow \bar{D}$ in \mathcal{D} , we prove that the morphisms $e_i \otimes e_i$ have the wide pushout $e \otimes e = (\bar{e}_i \otimes \bar{e}_i) \cdot (e_i \otimes e_i)$ in \mathcal{D} . Indeed, given a compatible family

$$c : D \otimes D \rightarrow C \text{ and } c_i : D_i \otimes D_i \rightarrow C \text{ with } c = c_i \cdot (e_i \otimes e_i),$$

the morphisms $e_i \otimes D$, $i \in I$, have the compatible family formed by c and $c_i \cdot (D_i \otimes e_i)$. Since $- \otimes D$ is a left adjoint and thus preserves colimits, the wide pushout of $e_i \otimes D$ is formed by $e \otimes D$ and $\bar{e}_i \otimes D$. Consequently there is unique morphism

$$\tilde{c} : \bar{D} \otimes D \rightarrow C$$

such that the following squares commute for all $i \in I$:

$$\begin{array}{ccc} D_i \otimes D & \xrightarrow{\bar{e}_i \otimes D} & \bar{D} \otimes D \\ D_i \otimes e_i \downarrow & & \downarrow \tilde{c} \\ D_i \otimes D_i & \xrightarrow{c_i} & C \end{array}$$

Since by assumption $\bar{e}_i \in \mathcal{E}$, the stability of \mathcal{D} gives a unique morphism \tilde{c}_i making the following diagram commutative:

$$\begin{array}{ccccc} & & D_i \otimes D & & \\ & \swarrow D_i \otimes e_i & & \searrow \bar{e}_i \otimes D & \\ D_i \otimes D_i & & & & \bar{D} \otimes D \\ & \searrow \bar{e}_i \otimes D_i & & \swarrow \bar{D} \otimes e_i & \\ & & \bar{D} \otimes D_i & & \\ & \swarrow c_i & \downarrow \tilde{c}_i & \searrow \tilde{c} & \\ & & C & & \end{array}$$

Thus \tilde{c} and \tilde{c}_i form a cocone of the family $\bar{D} \otimes e_i$. Since the latter family has the wide pushout formed by $\bar{D} \otimes e$ and $\bar{D} \otimes \bar{e}_i$ (using that the left adjoint $\bar{D} \otimes -$ preserves colimits), we obtain a unique morphism $\bar{c} : \bar{D} \otimes \bar{D} \rightarrow C$ for which the following triangle commutes:

$$\begin{array}{ccc} \bar{D} \otimes D_i & \xrightarrow{\bar{D} \otimes \bar{e}_i} & \bar{D} \otimes \bar{D} \\ & \searrow \tilde{c}_i & \downarrow \bar{c} \\ & & C \end{array}$$

Combined with the definition of \tilde{c}_i , we get

$$c_i = \tilde{c}_i \cdot (\bar{e}_i \otimes D_i) = \bar{c} \cdot (\bar{D} \otimes \bar{e}_i) \cdot (\bar{e}_i \otimes D_i) = \bar{c} \cdot (\bar{e}_i \otimes \bar{e}_i)$$

for all $i \in I$, as desired.

The uniqueness of \bar{c} follows from the fact that the morphisms $\bar{e}_i \otimes \bar{e}_i$ are epimorphisms (using that $\bar{D} \otimes -$ and $- \otimes D_i$ are left-adjoints and hence preserve epimorphisms).

- (2) Suppose that \mathcal{D} -monoid structures are given such that the above morphisms e_i are \mathcal{D} -monoid morphisms

$$e_i : (D, \mu, \eta) \rightarrow (D_i, \mu_i, \eta_i)$$

for all $i \in I$. We define a \mathcal{D} -monoid structure $(\bar{D}, \bar{\mu}, \bar{\eta})$ such that all \bar{e}_i are \mathcal{D} -monoid morphisms. For the unit $\bar{\eta}$ put

$$\bar{\eta} \equiv I \xrightarrow{\eta} D \xrightarrow{e} \bar{D}.$$

Next, observe that the morphisms $e_i \otimes e_i$ have a compatible cocone consisting of $e \cdot \mu$ and $\bar{e}_i \cdot \mu_i$ ($i \in I$). Hence by (1) there exists a unique $\bar{\mu} : \bar{D} \otimes \bar{D} \rightarrow \bar{D}$ for which the following squares commute:

$$\begin{array}{ccc} D \otimes D & \xrightarrow{\mu} & D \\ e \otimes e \downarrow & & \downarrow e \\ \bar{D} \otimes \bar{D} & \xrightarrow{\bar{\mu}} & \bar{D} \end{array} \quad \begin{array}{ccc} D_i \otimes D_i & \xrightarrow{\mu_i} & D_i \\ \bar{e}_i \otimes \bar{e}_i \downarrow & & \downarrow \bar{e}_i \\ \bar{D} \otimes \bar{D} & \xrightarrow{\bar{\mu}} & \bar{D} \end{array}$$

We only need to prove that $(\bar{D}, \bar{\mu}, \bar{\eta})$ is a \mathcal{D} -monoid. Then the definitions of μ and η immediately imply that e and \bar{e}_i are \mathcal{D} -monoid morphisms, and the verification that they form the wide pushout in $\mathbf{Mon}(D)$ is trivial.

Unit laws. Due to symmetry we only verify the left unit law $\bar{\mu} \cdot (\bar{\eta} \otimes \bar{D}) = \lambda_{\bar{D}}$, where $\lambda_{\bar{D}} : I \otimes \bar{D} \cong \bar{D}$ is the left unit isomorphism. Consider the following diagram:

$$\begin{array}{ccccc} I \otimes D & \xrightarrow{\eta \otimes D} & & & D \otimes D \\ & \searrow I \otimes e & & & \swarrow e \otimes e \\ & & I \otimes \bar{D} & \xrightarrow{\bar{\eta} \otimes \bar{D}} & \bar{D} \otimes \bar{D} \\ & \searrow \lambda_D & \searrow \lambda_{\bar{D}} & & \swarrow \mu \\ & & & & \bar{D} \\ & & & & \uparrow e \\ & & & & D \end{array}$$

The outside triangle is the unit law of (D, μ, η) . The upper part commutes due to $\bar{\eta} = e \cdot \eta$, the left-hand one is the naturality of λ , and the right-hand one is the definition of $\bar{\mu}$. Thus the inner triangle commutes when precomposed with $I \otimes e$. Since the latter is an epimorphism, the proof is complete.

Associative law. Consider the following diagram, where α denotes the associativity isomorphism:

$$\begin{array}{ccccc}
 (D \otimes D) \otimes D & \xrightarrow{\alpha} & D \otimes (D \otimes D) & \xrightarrow{D \otimes \mu} & D \otimes D \\
 \downarrow \mu \otimes I & \searrow (e \otimes e) \otimes e & \downarrow e \otimes (e \otimes e) & \swarrow e \otimes e & \downarrow \mu \\
 & (\bar{D} \otimes \bar{D}) \otimes \bar{D} & \xrightarrow{\alpha} & \bar{D} \otimes (\bar{D} \otimes \bar{D}) & \xrightarrow{\bar{D} \otimes \bar{\mu}} & \bar{D} \otimes \bar{D} \\
 & \downarrow \bar{\mu} \otimes \bar{D} & & & \downarrow \bar{\mu} \\
 & \bar{D} \otimes \bar{D} & \xrightarrow{\bar{\mu}} & \bar{D} & \\
 \downarrow e \otimes e & \swarrow e \otimes e & & & \downarrow e \\
 D \otimes D & \xrightarrow{\mu} & D & & D
 \end{array}$$

The outside is the associativity of (D, μ, η) . All the inner parts except for the (desired) inner square commute: for the upper left-hand one use the naturality of α , and all other parts follow from the definition of $\bar{\mu}$. Consequently the inner square commutes, as it commutes when precomposed with the epimorphism $(e \otimes e) \otimes e$. \square

Corollary 3.10. *Let \mathcal{D} satisfy the assumptions of the previous theorem, and suppose moreover that \mathcal{D} has wide pushouts of \mathcal{E} -morphisms. Then every language $L : X^{\otimes} \rightarrow Y$ has a syntactic \mathcal{D} -monoid.*

Proof. Let $e_i : X^{\otimes} \rightarrow M_i$ ($i \in I$) be the family of all X -generated \mathcal{D} -monoids recognizing L . Note that $I \neq \emptyset$ since the identity morphism of X^{\otimes} trivially recognizes L . Form the wide pushout $e : X^{\otimes} \rightarrow M$ and $\bar{e}_i : M_i \rightarrow M$ of all e_i 's in \mathcal{D} . By Theorem 3.9 there is a unique \mathcal{D} -monoid structure on M making e and \bar{e}_i a wide pushout in $\mathbf{Mon}(\mathcal{D})$. We claim that $e : X^{\otimes} \rightarrow M$ is the syntactic \mathcal{D} -monoid of L . To this end we verify the universal property of Definition 3.6:

(a) Since e_i recognizes L , there exists a morphism $f_i : M_i \rightarrow Y$ in \mathcal{D} with $L = f_i \cdot e_i$ for each $i \in I$. Hence the morphisms L and f_i form a compatible family, so there is a unique morphism $f : M \rightarrow Y$ in \mathcal{D} with $f_i = f \cdot \bar{e}_i$ for all i . Choosing an arbitrary $i \in I$, it follows that

$$f \cdot e = f \cdot \bar{e}_i \cdot e_i = f_i \cdot e_i = L,$$

i.e. e recognizes L via f .

(b) By construction, for every X -generated \mathcal{D} -monoid $e_i : X^{\otimes} \rightarrow M_i$ recognizing L we have the \mathcal{D} -monoid morphism $\bar{e}_i : M_i \rightarrow M$ with $\bar{e}_i \cdot e_i = e$. \square

Corollary 3.11. *Every language $L : X^{\otimes} \rightarrow Y$ in a variety \mathcal{D} of algebras or ordered algebras admits a syntactic \mathcal{D} -monoid.*

We shall now show that in the situation of the above corollary the syntactic \mathcal{D} -monoid of a language L admits a more concrete construction via the *syntactic congruence* of L . We first consider the case of varieties of algebras (Section 3.2) and then turn to varieties of ordered algebras (Section 3.3).

3.2. Syntactic congruences for varieties of algebras. In this subsection, let \mathcal{D} be a variety of algebras.

Remark 3.12. (1) Recall that a *congruence* on a Σ -algebra A is an equivalence relation \equiv on A that forms a subalgebra of $A \times A$. We denote by A/\equiv the quotient algebra modulo \equiv .

(2) For any homomorphism $h : A \rightarrow B$ of Σ -algebras, the *kernel congruence* of h is the congruence on A defined by

$$a \equiv_h a' \quad \text{iff} \quad h(a) = h(a').$$

(3) We will frequently use the following *homomorphism theorem*: given homomorphisms of Σ -algebras $e : A \rightarrow B$ and $h : A \rightarrow C$, where e is surjective, there exists a homomorphism $h' : B \rightarrow C$ with $h = h' \cdot e$ iff, for all $a, a' \in A$,

$$e(a) = e(a') \quad \text{implies} \quad h(a) = h(a').$$

Definition 3.13. Let \mathcal{D} be a variety of algebras. The *syntactic congruence* of a language $L : X^\otimes \rightarrow Y$ is the relation \equiv_L on $|X^\otimes|$ defined by

$$u \equiv_L v \quad \text{iff} \quad \forall x, y \in |X^\otimes| : L(x \bullet u \bullet y) = L(x \bullet v \bullet y). \quad (3.1)$$

Theorem 3.14. \equiv_L is a congruence on the free \mathcal{D} -monoid X^\otimes , and

$$\text{Syn}(L) = X^\otimes / \equiv_L.$$

Proof. (a) Clearly \equiv_L forms an equivalence relation on X^\otimes . To show that it is congruence, first observe that \equiv_L is a subobject of $X^\otimes \times X^\otimes$ in \mathcal{D} . Indeed, we have $\equiv_L = \bigcap K_{x,y}$, where for fixed $x, y \in X^\otimes$ the object $K_{x,y}$ is the kernel of the \mathcal{D} -morphism $X^\otimes \xrightarrow{x \bullet -} X^\otimes \xrightarrow{- \bullet y} X^\otimes \xrightarrow{L} Y$. Moreover, \equiv_L is closed under the monoid multiplication of $X^\otimes \times X^\otimes$: given $u \equiv_L v$ and $u' \equiv_L v'$ we have for all $x, y \in X^\otimes$ that

$$\begin{aligned} L(x \bullet u \bullet u' \bullet y) &= L(x \bullet v \bullet u' \bullet y) && (y := u' \bullet y \text{ in (3.1)}) \\ &= L(x \bullet v \bullet v' \bullet y) && (x := x \bullet v \text{ in (3.1)}) \end{aligned}$$

Hence $u \bullet u' \equiv_L v \bullet v'$ and therefore \equiv_L is a \mathcal{D} -submonoid of $X^\otimes \times X^\otimes$, i.e. a congruence of X^\otimes .

(b) Denote by $e_L : X^\otimes \rightarrow X^\otimes / \equiv_L$ the projection. Then $e_L(u) = e_L(v)$ (i.e. $u \equiv_L v$) implies $L(u) = L(v)$ by putting $x = y = 1$ in (3.1), where 1 is the unit of the monoid X^\otimes . Hence the homomorphism theorem yields a morphism $f_L : X^\otimes / \equiv_L \rightarrow Y$ in \mathcal{D} with $f_L \cdot e_L = L$, i.e. e_L recognizes L via f_L .

(c) It remains to verify the universal property of Definition 3.5. Let $e : X^\otimes \rightarrow M$ be a surjective \mathcal{D} -monoid morphism recognizing L via $f : M \rightarrow Y$. To construct the morphism $h : M \rightarrow X^\otimes / \equiv_L$, we again use the homomorphism theorem. Thus let $u, v \in X^\otimes$ with $e(u) = e(v)$. Then, for all $x, y \in X^\otimes$,

$$\begin{aligned} L(x \bullet u \bullet y) &= f(e(x \bullet u \bullet y)) && (L = f \cdot e) \\ &= f(e(x)e(u)e(y)) && (e \text{ preserves } \bullet) \\ &= f(e(x)e(v)e(y)) && (e(u) = e(v)) \\ &= f(e(x \bullet v \bullet y)) && (e \text{ preserves } \bullet) \\ &= L(x \bullet v \bullet y) && (L = f \cdot e) \end{aligned}$$

Hence $u \equiv_L v$, i.e. $e_L(u) = e_L(v)$. Thus the homomorphism theorem yields the desired \mathcal{D} -monoid morphism $h : M \rightarrow X^{\otimes} / \equiv_L$ with $h \cdot e = e_L$. \square

Example 3.15. Using the notation of Example 2.29 we obtain the following syntactic algebras:

- (1) In **Set** with $Y = \{0, 1\}$, the syntactic monoid of a language $L \subseteq X^*$ is the quotient monoid X^* / \equiv_L , where for $u, v \in X^*$,

$$u \equiv_L v \quad \text{iff} \quad \text{for all } x, y \in X^* : xuy \in L \iff xvy \in L.$$

This construction is due to Rabin and Scott [19].

- (2) In **Set** $_{\perp}$ with $X = X_0 + \{\perp\}$ and $Y = \{\perp, 1\}$ the *syntactic monoid with zero* of a language $L_0 \subseteq X_0^*$ is $(X_0^* + \{\perp\}) / \equiv_L$ where for $u, v \in X_0^* + \{\perp\}$,

$$u \equiv_L v \quad \text{iff} \quad \text{for all } x, y \in X_0^* : xuy \in L_0 \iff xvy \in L_0.$$

The zero element is the congruence class of \perp .

- (3) In **Inv** with $X = X_0 + \widetilde{X}_0$ and $Y = \{0, 1\}$ the *syntactic involution monoid* of a language $L_0 \subseteq X_0^*$ is the quotient of $X_0 + \widetilde{X}_0^*$ modulo the congruence \equiv_L defined for words $u, v \in X_0^*$ as follows:

- (i) $u \equiv_L v$ iff $\widetilde{u} \equiv_L \widetilde{v}$ iff for all $x, y \in X_0^* : xuy \in L_0 \iff xvy \in L_0$;
(ii) $u \equiv_L \widetilde{v}$ iff $\widetilde{u} \equiv_L v$ iff for all $x, y \in X_0^* : xuy \in L_0 \iff xvy \notin L_0$.

- (4) In **Mod**(\mathbb{S}) with $X = \Psi X_0$ and $Y = \mathbb{S}$ the *syntactic associative \mathbb{S} -algebra* of a weighted language $L_0 : X_0^* \rightarrow \mathbb{S}$ is the quotient of $\mathbb{S}[X_0]$ modulo the congruence defined for $U, V \in \mathbb{S}[X_0]$ as follows:

$$U \equiv_L V \quad \text{iff} \quad \text{for all } x, y \in X_0^* : L(xUy) = L(xVy) \tag{3.2}$$

Indeed, since $L : \mathbb{S}[X_0] \rightarrow \mathbb{S}$ is linear, (3.2) implies $L(PUQ) = L(PVQ)$ for all $P, Q \in \mathbb{S}[X_0]$, which is the syntactic congruence of Definition 3.13.

- (5) In particular, for $\mathcal{D} = \mathbf{JSL}_0$ with $X = \mathcal{P}_f X_0$ and $Y = \{0, 1\}$, we get the *syntactic (idempotent) semiring* of a language $L_0 \subseteq X_0^*$ introduced by Polák [18]: it is the quotient $\mathcal{P}_f X_0^* / \equiv_L$ where for $U, V \in \mathcal{P}_f X_0^*$ we have

$$U \equiv_L V \quad \text{iff} \quad \text{for all } x, y \in X_0^* : (xUy) \cap L_0 \neq \emptyset \iff xVy \cap L_0 \neq \emptyset.$$

- (6) For $\mathcal{D} = \mathbf{Vec}(\mathbb{K})$ with $X = \Psi X_0$ and $Y = \mathbb{K}$, the *syntactic \mathbb{K} -algebra* of a \mathbb{K} -weighted language $L_0 : X_0^* \rightarrow \mathbb{K}$ is the quotient $\mathbb{K}[X_0] / I$ of the \mathbb{K} -algebra of finite weighted languages modulo the ideal

$$I = \{V \in \mathbb{K}[X_0] \mid \text{for all } x, y \in X_0^* : L(xVy) = 0\}.$$

Indeed, the congruence this ideal I generates ($U \equiv_L V$ iff $U - V \in I$) is precisely (3.2). Syntactic \mathbb{K} -algebras were studied by Reutenauer [20].

- (7) Analogously, for $\mathcal{D} = \mathbf{Ab}$ with $X = \Psi X_0$ and $Y = \mathbb{Z}$, the *syntactic ring* of a \mathbb{Z} -weighted language $L_0 : X_0^* \rightarrow \mathbb{Z}$ is the quotient $\mathbb{Z}[X_0] / I$, where I is the ideal of all $V \in \mathbb{Z}[X_0]$ with $L(xVy) = 0$ for all $x, y \in X_0^*$.

3.3. Syntactic congruences for ordered algebras. Next, we consider the case where \mathcal{D} is a variety of ordered algebras.

Remark 3.16. (1) A *congruence* on an ordered Σ -algebra (A, \leq) is a preorder \leq on A such that $a \leq a'$ implies $a \leq a'$, and all Σ -operations of A are monotone w.r.t. \leq (equivalently, \leq forms a subalgebra of $A \times A$). Then $\leq \cap \geq$ is a congruence in the unordered sense, and the quotient algebra $e : A \twoheadrightarrow A/(\leq \cap \geq)$ forms an ordered algebra with partial order induced by e :

$$e(u) \leq' e(v) \quad \text{iff} \quad u \leq v.$$

This ordered algebra is called the *quotient* of A modulo \leq , and is denoted by A/\leq .

(2) For any homomorphism $h : A \rightarrow B$ between ordered Σ -algebras, the *kernel congruence* of h is the congruence on A defined by

$$a \leq_h a' \quad \text{iff} \quad h(a) \leq h(a').$$

(3) The *homomorphism theorem* has the following version for ordered algebras, see e.g. [17, Proposition 1.1]: given homomorphisms of ordered Σ -algebras $e : A \twoheadrightarrow B$ and $h : A \rightarrow C$, where e is surjective, there exists a homomorphism $h' : B \rightarrow C$ with $h = h' \cdot e$ iff, for all $a, a' \in A$,

$$e(a) \leq e(a') \quad \text{implies} \quad h(a) \leq h(a').$$

Definition 3.17. Let \mathcal{D} be a variety of ordered algebras. The *syntactic congruence* of a language $L : X^\otimes \rightarrow Y$ is the relation \leq_L on $|X^\otimes|$ defined by

$$u \leq_L v \quad \text{iff} \quad \forall x, y \in |X^\otimes| : L(x \bullet u \bullet y) \leq L(x \bullet v \bullet y) \quad (3.3)$$

Theorem 3.18. \leq_L is a congruence of the free \mathcal{D} -monoid X^\otimes , and

$$\text{Syn}(L) = X^\otimes / \leq_L.$$

Proof. Clearly \leq_L is a preorder. Let us verify the conditions of Remark 3.16(1). First, if $u \leq v$ in X^\otimes , then $x \bullet u \bullet y \leq x \bullet v \bullet y$ for all $x, y \in X^\otimes$ because \bullet is monotone. Since also L is monotone, we conclude $L(x \bullet u \bullet y) \leq L(x \bullet v \bullet y)$ in Y for all x, y , i.e. $u \leq_L v$. Next, observe that \leq_L is a subobject of $X^\otimes \times X^\otimes$ in \mathcal{D} , namely $\leq_L = \bigcap \leq_{x,y}$, where for fixed $x, y \in X^\otimes$ the object $\leq_{x,y}$ is the kernel congruence of the \mathcal{D} -morphism $X^\otimes \xrightarrow{x \bullet -} X^\otimes \xrightarrow{- \bullet y} X^\otimes \xrightarrow{L} Y$. Moreover, the monoid multiplication \bullet is monotone w.r.t. \leq_L : given $u \leq_L v$ and $u' \leq_L v'$ we have for all $x, y \in X^\otimes$ that

$$L(x \bullet u \bullet u' \bullet y) \leq L(x \bullet v \bullet u' \bullet y) \leq L(x \bullet v \bullet v' \bullet y).$$

Hence $u \bullet u' \leq_L v \bullet v'$, and therefore \leq_L is a \mathcal{D} -submonoid of $X^\otimes \times X^\otimes$, as required.

The proof that $\text{Syn}(L) = X^\otimes / \leq_L$ is completely analogous to that of Proposition 3.14: replace equations by inequations, and use the homomorphism theorem for ordered algebras to construct the morphisms f_L and h . \square

Example 3.19. In $\mathcal{D} = \mathbf{Pos}$ with $Y = \{0 < 1\}$, the *syntactic ordered monoid* of a language $L \subseteq X^*$ is the ordered quotient monoid X^* / \leq_L where for $u, v \in X^*$,

$$u \leq_L v \quad \text{iff} \quad \text{for all } x, y \in X^* : xuy \in L \Rightarrow xvy \in L.$$

This construction is due to Pin [16].

4. TRANSITION \mathcal{D} -MONOIDS

In this section we present another construction of the syntactic \mathcal{D} -monoid of a language: it is the transition \mathcal{D} -monoid of the minimal \mathcal{D} -automaton for this language. We continue to work under the Assumptions 2.7. Recall from Remark 2.22 the \mathcal{D} -monoid $[Q, Q]$ and the \mathcal{D} -monoid morphism $(\lambda\delta)^+ : X^\otimes \rightarrow [Q, Q]$.

Definition 4.1. The *transition \mathcal{D} -monoid* $\mathbb{T}(Q)$ of an F -algebra (Q, δ, i) is the image of the \mathcal{D} -monoid morphism $(\lambda\delta)^+ : X^\otimes \rightarrow [Q, Q]$:

$$\begin{array}{ccc} X^\otimes & \xrightarrow{(\lambda\delta)^+} & [Q, Q] \\ & \searrow e_{\mathbb{T}(Q)} & \nearrow m_{\mathbb{T}(Q)} \\ & \mathbb{T}(Q) & \end{array}$$

- Example 4.2.** (1) In **Set** or **Pos** the (ordered) transition monoid of an F -algebra Q , i.e. an (ordered) automaton without final states, is the (ordered) monoid of all extended transition maps $\delta_w = \delta_{a_n} \cdot \dots \cdot \delta_{a_1} : Q \rightarrow Q$ for $w = a_1 \cdot \dots \cdot a_n \in X^*$. The unit is $\text{id}_Q = \delta_\varepsilon$ and the monoid multiplication is composition.
- (2) In **Set** $_\perp$ with $X = X_0 + \{\perp\}$ (the setting for partial automata) this is completely analogous, except that we add the constant endomap of Q with value \perp .
- (3) In **Inv** with $X = X_0 + \widetilde{X}_0$ we get the involution monoid of all δ_w and $\widetilde{\delta}_w$. Again the unit is δ_ε , and the multiplication is determined by composition plus the equations $x\widetilde{y} = \widetilde{xy} = \widetilde{xy}$.
- (4) In **JSL** $_0$ with $X = \mathcal{P}_f X_0$ the transition semiring consists of all finite joins of extended transitions, i.e. all semilattice homomorphisms of the form $\delta_{w_1} \vee \dots \vee \delta_{w_n}$ for $\{w_1, \dots, w_n\} \in \mathcal{P}_f X_0^*$. The transition semiring was introduced by Polák [18].
- (5) In **Mod**(\mathbb{S}) with $X = \Psi X_0$ the associative transition algebra consists of all linear maps of the form $\sum_{i=1}^n s_i \delta_{w_i}$ with $s_i \in \mathbb{S}$ and $w_i \in X_0^*$.

Recall from Definition 2.16 that a \mathcal{D} -automaton is an F -algebra Q together with an output morphism $f : Q \rightarrow Y$. Hence we can speak of the transition \mathcal{D} -monoid of a \mathcal{D} -automaton.

Proposition 4.3. The language accepted by a \mathcal{D} -automaton (Q, δ, f, i) is recognized by the \mathcal{D} -monoid morphism $e_{\mathbb{T}(Q)} : X^\otimes \rightarrow \mathbb{T}(Q)$.

Proof. Let (Q, δ, i, f) be a \mathcal{D} -automaton. By definition it accepts the language $L_Q = (X^\otimes \xrightarrow{e_Q} Q \xrightarrow{f} Y)$ where e_Q is the unique F -algebra homomorphism. Consider the morphism that evaluates any endomorphism of Q at the initial state:

$$\text{ev}_i = ([Q, Q] \cong [Q, Q] \otimes I \xrightarrow{[Q, Q] \otimes i} [Q, Q] \otimes Q \xrightarrow{\text{ev}} Q).$$

Now let

$$f_{\mathbb{T}(Q)} = (\mathbb{T}(Q) \xrightarrow{m_{\mathbb{T}(Q)}} [Q, Q] \xrightarrow{\text{ev}_i} Q \xrightarrow{f} Y).$$

With this morphism $\mathsf{T}(Q)$ recognizes L ; indeed, using the right unit isomorphism $\iota_Z : Z \rightarrow Z \otimes I$ we compute:

$$\begin{aligned}
L_Q &= f \cdot e_Q \\
&= f \cdot \mathbf{ev} \cdot ((\lambda\delta)^+ \otimes i) \cdot \iota_{X^\otimes} && \text{(Remark 2.22)} \\
&= f \cdot \mathbf{ev}_i \cdot \iota_{[Q,Q]}^{-1} \cdot ((\lambda\delta)^+ \otimes I) \cdot \iota_{X^\otimes} && \text{(def. of } \mathbf{ev}_i) \\
&= f \cdot \mathbf{ev}_i \cdot (\lambda\delta)^+ && \text{(naturality of } \iota) \\
&= f \cdot \mathbf{ev}_i \cdot m_{\mathsf{T}(Q)} \cdot e_{\mathsf{T}(Q)} && \text{(Definition 4.1)} \\
&= f_{\mathsf{T}(Q)} \cdot e_{\mathsf{T}(Q)} && \text{(def. of } f_{\mathsf{T}(Q)})
\end{aligned}$$

This completes the proof. \square

Definition 4.4. A \mathcal{D} -automaton (Q, δ, i, f) is called *minimal* iff it is

- (a) *reachable*: the unique F -algebra homomorphism $X^\otimes \rightarrow Q$ is surjective;
- (b) *simple*: the unique T -coalgebra homomorphism $Q \rightarrow [X^\otimes, Y]$ is injective.

Theorem 4.5 (Goguen [12]). *Every language $L : X^\otimes \rightarrow Y$ is accepted by a minimal \mathcal{D} -automaton $\text{Min}(L)$, unique up to isomorphism. Given any reachable automaton Q accepting L , there is a unique surjective automata homomorphism from Q into $\text{Min}(L)$.*

This leads to the announced construction of syntactic \mathcal{D} -monoids via transition \mathcal{D} -monoids. The case $\mathcal{D} = \mathbf{Set}$ and \mathbf{Pos} are standard results of algebraic automata theory (see e.g. Pin [15]), and the case $\mathcal{D} = \mathbf{JSL}_0$ is due to Polák [18]. For the other instances in Example 4.2 this appears to be new.

Remark 4.6. Recall that in our algebraic setting the tensor unit I is $\Psi 1$. Hence elements $a \in A$ of a \mathcal{D} -object are in 1-1-correspondence with morphisms $a : I \rightarrow A$; we shall henceforth not distinguish these.

Theorem 4.7. *The syntactic \mathcal{D} -monoid of a language $L : X^\otimes \rightarrow Y$ is isomorphic to the transition \mathcal{D} -monoid of its minimal \mathcal{D} -automaton:*

$$\text{Syn}(L) \cong \mathsf{T}(\text{Min}(L)).$$

Proof. Let $\text{Min}(L) = (Q, \delta, i, f)$, and write $\delta_x : Q \rightarrow Q$ for $e_{\mathsf{T}(Q)}(x)$ ($x \in X^\otimes$). Note that $\delta_{x \bullet y} = \delta_y \cdot \delta_x$ for all $x, y \in X^\otimes$ since $e_{\mathsf{T}(Q)}$ is a \mathcal{D} -monoid morphism. Recall also that the unique F -algebra homomorphism $e_Q : X^\otimes \rightarrow Q$ assigns to $x \in X^\otimes$ the element $\delta_x \cdot i : I \rightarrow Q$ (see Remark 2.22(2)), and the unique T -coalgebra homomorphism $m_Q : Q \rightarrow [X^\otimes, Y]$ assigns to a state $q : I \rightarrow Q$ the language $x \mapsto f \cdot \delta_x \cdot q$ (see Remark 2.27). It suffices to show that the kernel congruence of $e_{\mathsf{T}(Q)}$ coincides with the syntactic congruence of L . If \mathcal{D} is a variety of algebras, this requires to prove that for all $u, v \in |X^\otimes|$ one has

$$\delta_u = \delta_v \quad \text{iff} \quad \forall x, y \in X^\otimes : L(x \bullet u \bullet y) = L(x \bullet v \bullet y).$$

To see this, we compute

$$\begin{aligned}
\delta_u = \delta_v &\Leftrightarrow \forall x : \delta_u \cdot e_Q(x) = \delta_v \cdot e_Q(x) && (e_Q \text{ surjective}) \\
&\Leftrightarrow \forall x : \delta_u \cdot \delta_x \cdot i = \delta_v \cdot \delta_x \cdot i && (\text{def. } e_Q) \\
&\Leftrightarrow \forall x : m_Q \cdot \delta_u \cdot \delta_x \cdot i = m_Q \cdot \delta_v \cdot \delta_x \cdot i && (m_Q \text{ injective}) \\
&\Leftrightarrow \forall x, y : f \cdot \delta_y \cdot \delta_u \cdot \delta_x \cdot i = f \cdot \delta_y \cdot \delta_v \cdot \delta_x \cdot i && (\text{def. } m_Q) \\
&\Leftrightarrow \forall x, y : f \cdot \delta_{x \bullet u \bullet y} \cdot i = f \cdot \delta_{x \bullet v \bullet y} \cdot i && (\text{def. } \delta_{(-)}) \\
&\Leftrightarrow \forall x, y : f \cdot e_Q(x \bullet u \bullet y) = f \cdot e_Q(x \bullet v \bullet y) && (\text{def. } e_Q) \\
&\Leftrightarrow \forall x, y : L(x \bullet u \bullet y) = L(x \bullet v \bullet y) && (L = L_Q)
\end{aligned}$$

The case where \mathcal{D} is a variety of ordered algebras is completely analogous: just replace equations by inequations. \square

5. \mathcal{D} -REGULAR LANGUAGES

Our results so far apply to arbitrary languages in \mathcal{D} . In the present section we focus on *regular languages*, which in $\mathcal{D} = \mathbf{Set}$ are the languages accepted by finite automata, or equivalently the languages recognized by finite monoids. For arbitrary \mathcal{D} the role of finite sets is taken over by finitely presentable objects. Recall that an object D of \mathcal{D} is *finitely presentable* if the hom-functor $\mathcal{D}(D, -) : \mathcal{D} \rightarrow \mathbf{Set}$ preserves filtered colimits. Equivalently, D can be presented with finitely many generators and relations. For example, the tensor unit $I = \Psi 1$ is finitely presentable, since it is presented with one generator and no relations.

Definition 5.1. A language $L : X^* \rightarrow Y$ is called *\mathcal{D} -regular* if it is accepted by some \mathcal{D} -automaton with a finitely presentable object of states.

To work with this definition, we need the following

Assumptions 5.2. We assume that the full subcategory \mathcal{D}_f of finitely presentable objects of \mathcal{D} is closed under subobjects, quotients and finite products.

Example 5.3. (1) Recall that a variety is *locally finite* if all finitely presentable algebras (equivalently all finitely generated free algebras) are finite. Every locally finite variety satisfies the above assumptions. This includes our examples \mathbf{Set} , \mathbf{Set}_\perp , \mathbf{Inv} and \mathbf{JSL}_0 .
(2) A semiring \mathbb{S} is called *Noetherian* if all submodules of finitely generated \mathbb{S} -modules are finitely generated. In this case, as shown in [10], the category $\mathbf{Mod}(\mathbb{S})$ satisfies our assumptions. Every field is Noetherian, as is every finitely generated commutative ring, so $\mathbf{Vec}(\mathbb{K})$ and $\mathbf{Ab} = \mathbf{Mod}(\mathbb{Z})$ are special instances.

Example 5.4. As we have mentioned already, for $\mathcal{D} = \mathbf{Set}$ the \mathcal{D} -regular languages are precisely the classical regular languages. The same is true for \mathbf{Set}_\perp , \mathbf{Pos} , \mathbf{Inv} and \mathbf{JSL}_0 . For $\mathcal{D} = \mathbf{Mod}(\mathbb{S})$ for a Noetherian semiring \mathbb{S} it is known that the \mathcal{D} -regular languages are precisely the rational weighted languages, see e.g. [10].

Remark 5.5. (1) The functor $FQ = I + X \otimes Q$ preserves surjective homomorphisms. Therefore the factorization system of \mathcal{D} lifts to the category of F -algebras, that is, every F -algebra homomorphism factorizes into a surjective homomorphism followed by an injective (resp. order-reflecting) one. By the *reachable part* Q_r of a \mathcal{D} -automaton (Q, δ, f, i) we mean the image of the initial F -algebra homomorphism,

i.e. $e_Q = (X^\otimes \xrightarrow{e_r} Q_r \xrightarrow{m_r} Q)$. Putting $f_r := f \cdot m_r : Q_r \rightarrow Y$, the F -algebra Q_r becomes an automaton, and m_r an automata homomorphism. \square

(2) In the following automata, (co-)algebras and monoids with finitely presentable carrier are referred to as *fp-automata*, *fp-(co-)algebras* and *fp-monoids*, respectively.

Theorem 5.6. *For any language $L : X^\otimes \rightarrow Y$ the following statements are equivalent:*

- (a) L is \mathcal{D} -regular.
- (b) The minimal \mathcal{D} -automaton $\text{Min}(L)$ has a finitely presentable carrier.
- (c) L is recognized by some \mathcal{D} -monoid with a finitely presentable carrier.
- (d) The syntactic \mathcal{D} -monoid $\text{Syn}(L)$ has a finitely presentable carrier.

Proof. (a) \Leftrightarrow (b) follows from Theorem 4.5 and our assumption that \mathcal{D}_f is closed under subobjects and quotients. Similarly, (c) \Leftrightarrow (d) follows from the universal property of the syntactic monoid (see Definition 3.6) and again closure of \mathcal{D}_f under subobjects and quotients. (c) \Rightarrow (a) is a consequence of Lemma 3.5. To prove (a) \Rightarrow (c), let Q be any fp-automaton accepting L . Then by Proposition 4.3 the transition monoid $\mathbb{T}(Q) \mapsto [Q, Q]$ recognizes L , so by closedness of \mathcal{D}_f under subobjects it suffices to show that $[Q, Q]$ is a finitely presentable object of \mathcal{D} . Assuming that Q has n generators as an algebra of \mathcal{D} , the map $[Q, Q] \rightarrow Q^n$ defined by restriction to the set of generators is an injective (resp. order-reflecting) morphism of \mathcal{D} . Since \mathcal{D}_f is closed under subobjects and finite products, it follows that $[Q, Q]$ is finitely presentable. \square

Just as the collection of all languages is internalized by the final coalgebra $[X^\otimes, Y]$, see Proposition 2.26, we can internalize the regular languages by means of the *rational coalgebra*.

Definition 5.7. The *rational coalgebra* ϱT for T is the colimit (taken in the category of T -coalgebras and homomorphisms) of all T -coalgebras with finitely presentable carrier.

Example 5.8. In $\mathcal{D} = \mathbf{Set}$, the rational coalgebra is the subcoalgebra of the final coalgebra $\mathcal{P}X_0^*$ given by the set of all regular languages. Analogously for $\mathcal{D} = \mathbf{Pos}$ where the order is given by inclusion of languages. In general, ϱT always has the \mathcal{D} -regular languages as states:

Proposition 5.9. *There is a one-to-one correspondence between \mathcal{D} -regular languages and elements $I \rightarrow \varrho T$ of the rational coalgebra.*

Proof. We describe mutually inverse maps

$$(I \xrightarrow{x} \varrho T) \mapsto (X^\otimes \xrightarrow{L_x} Y) \quad \text{and} \quad (X^\otimes \xrightarrow{L} Y) \mapsto (I \xrightarrow{x_L} \varrho T)$$

between the elements of ϱT and the \mathcal{D} -regular languages. Let $h_Q : Q \rightarrow \varrho T$ be the injections of the colimit ϱT , where $Q = (Q, \delta_Q, f_Q)$ ranges over all fp-coalgebras. Note that this colimit is filtered because \mathcal{D}_f is closed under finite colimits in \mathcal{D} . Moreover, since colimits of coalgebras are formed in the underlying category, the morphisms h_Q also form a colimit cocone in \mathcal{D} . Given an element $I \xrightarrow{x} \varrho T$ of the rational coalgebra we define a \mathcal{D} -regular language $L_x : X^\otimes \rightarrow Y$ as follows: since $I = \Psi 1$ is finitely presentable, there exists an fp-coalgebra Q and a morphism $i_Q : I \rightarrow Q$ such that $x = h_Q \cdot i_Q$. For the F -algebra (Q, δ_Q, i_Q) we have the unique F -algebra homomorphism $e_Q : X^\otimes \rightarrow Q$, and we put $L_x := f_Q \cdot e_Q$.

We need to show that L_x is well-defined, that is, for any other factorization $x = h_{Q'} \cdot i_{Q'}$ we have $f_Q \cdot e_Q = f_{Q'} \cdot e_{Q'}$. Given such a factorization, since the morphisms h_Q form a filtered colimit cocone, there exists an fp-coalgebra $Q'' = (Q'', \delta_{Q''}, f_{Q''})$ and coalgebra homomorphisms $h_{Q''} : Q \rightarrow Q''$ and $h_{Q'Q''} : Q' \rightarrow Q''$ with $h_{Q''} \cdot i_Q = h_{Q'Q''} \cdot i_{Q'} =: i_{Q''}$.

Then for the F -algebra $(Q'', \delta_{Q''}, i_{Q''})$ we have the unique homomorphism $e_{Q''} : X^{\otimes} \rightarrow Q''$. Moreover, $h_{QQ''}$ and $h_{Q'Q''}$ are also homomorphisms of F -algebras. It follows that

$$\begin{aligned} f_Q \cdot e_Q &= f_{Q''} \cdot h_{QQ''} \cdot e_Q && (h_{QQ''} \text{ coalgebra homomorphism}) \\ &= f_{Q''} \cdot e_{Q''} && (h_{Q'Q''} F\text{-algebra hom.}, X^{\otimes} \text{ initial}) \end{aligned}$$

and analogously $f_{Q'} \cdot e_{Q'} = f_{Q''} \cdot e_{Q''}$. Hence $f_Q \cdot e_Q = f_{Q'} \cdot e_{Q'}$, as claimed.

Conversely, let $L : X^{\otimes} \rightarrow Y$ be a \mathcal{D} -regular language. Then there exists an fp-automaton (Q, δ_Q, i_Q, f_Q) with $L = f_Q \cdot e_Q$. Put $x_L := h_Q \cdot i_Q : I \rightarrow \varrho T$. To prove the well-definedness of x_L , consider the automata homomorphisms

$$Q \xleftarrow{m} \langle Q_r - \xrightarrow{e} \gg \text{Min}(L)$$

of Theorem 4.5. Then

$$\begin{aligned} h_Q \cdot i_Q &= h_Q \cdot m \cdot i_{Q_r} && (m \text{ algebra hom.}) \\ &= h_{Q_r} \cdot i_{Q_r} && ((h_Q) \text{ cocone and } m \text{ coalgebra hom.}) \\ &= h_{\text{Min}(L)} \cdot e \cdot i_{Q_r} && ((h_Q) \text{ cocone and } e \text{ coalgebra hom.}) \\ &= h_{\text{Min}(L)} \cdot i_{\text{Min}(L)} && (e \text{ algebra hom.}) \end{aligned}$$

Hence $x_L = h_Q \cdot i_Q$ is independent of the choice of Q . It now follows immediately from the definitions that $x \mapsto L_x$ and $L \mapsto x_L$ are mutually inverse and hence define the desired bijective correspondence. \square

6. DUAL CHARACTERIZATION OF SYNTACTIC \mathcal{D} -MONOIDS

We conclude this paper with a dual approach to syntactic monoids. This section is largely based on results from our papers [1, 3] where a categorical generalization of Eilenberg's variety theorem was proved. We work with the following

Assumptions 6.1. From now on \mathcal{D} is a locally finite commutative variety of algebras or ordered algebras (cf. Example 5.3) whose epimorphisms are surjective. Moreover, we assume that there is another locally finite variety \mathcal{C} of algebras such that the full subcategories \mathcal{C}_f and \mathcal{D}_f of finite algebras are dually equivalent. (Two such varieties \mathcal{C} and \mathcal{D} are called *predual*.)

The action of the equivalence functor $\mathcal{C}_f^{op} \xrightarrow{\cong} \mathcal{D}_f$ on objects and morphisms is denoted by $Q \mapsto \widehat{Q}$ and $f \mapsto \widehat{f}$. Letting $\bar{I} \in \mathcal{C}_f$ denote the free one-generated object of \mathcal{C} , we choose the output object $Y \in \mathcal{D}_f$ to be the dual object of \bar{I} . Moreover, let $\bar{Y} \in \mathcal{C}_f$ be the dual object of $I = \Psi 1 \in \mathcal{D}_f$, the free one-generated object of \mathcal{D} . Finally, we put $X = \Psi X_0$ for a finite alphabet X_0 . Note that the underlying sets of \bar{Y} and Y are canonically isomorphic:

$$|\bar{Y}| \cong \mathcal{C}(\bar{I}, \bar{Y}) \cong \mathcal{D}(I, Y) \cong |Y|. \quad (6.1)$$

To simplify the presentation, we will assume in the following that \bar{Y} and Y have a two-element underlying set $\{0, 1\}$. This is, however, inessential; see Remark 6.13 at the end of this section.

Example 6.2. The categories \mathcal{C} and \mathcal{D} in the table below satisfy our assumptions.

\mathcal{C}	\mathcal{D}
BA	Set
DL₀₁	Pos
BR	Set_⊥
JSL₀	JSL₀
Vec(\mathbb{Z}_2)	Vec(\mathbb{Z}_2)

Here **BA**, **DL₀₁** and **BR** are the categories of boolean algebras, distributive lattices with 0 and 1, and non-unital boolean rings (i.e. rings without 1 satisfying the equation $x \cdot x = x$). The preduality of **BA** and **Set** is a restriction of the well-known Stone duality: the dual equivalence functor $\mathbf{BA}_f^{op} \xrightarrow{\cong} \mathbf{Set}_f$ assigns to a finite boolean algebra B the set $\text{At}(B)$ of its atoms, and to a homomorphism $h : A \rightarrow B$ the map $\text{At}(h) : \text{At}(B) \rightarrow \text{At}(A)$ sending $b \in \text{At}(B)$ to the unique atom $a \in \text{At}(A)$ with $ha \geq b$. Using a similar Stone-type duality, we proved in [3] that the the category **BR** of non-unital boolean rings is predual to **Set_⊥**. The preduality between **DL₀₁** and **Pos** is due to Birkhoff. The last two examples above correspond to the well-known self-duality of finite semilattices and finite-dimensional vector spaces, respectively. We refer to [3] for details.

On \mathcal{C} we consider the endofunctor $\bar{T}Q = \bar{Y} \times Q^{X_0}$, where Q^{X_0} denotes the $|X_0|$ -fold power of Q in \mathcal{C} . Its coalgebras are precisely deterministic automata in \mathcal{C} without an initial state, represented as triples (Q, γ_a, f) with transition morphisms $\gamma_a : Q \rightarrow Q$ ($a \in X_0$) and an output morphism $f : Q \rightarrow \bar{Y}$.

In the following we use the case $\mathcal{C} = \mathbf{BA}$ and $\mathcal{D} = \mathbf{Set}$ as our only running example. For details on the other examples see [1, 3]

Example 6.3. In $\mathcal{C} = \mathbf{BA}$ a \bar{T} -coalgebra is a deterministic automaton with a boolean algebra Q of states, transition maps γ_a which are boolean homomorphisms, and an output map $f : Q \rightarrow \{0, 1\}$ that specifies (via the preimage of 1) an ultrafilter $F \subseteq Q$ of final states.

The rational coalgebra $\varrho\bar{T}$ of \bar{T} (i.e. the colimit of all finite \bar{T} -coalgebras) has as states the regular languages over X_0 . The final state predicate $f : \varrho\bar{T} \rightarrow \bar{Y} = \{0, 1\}$ sends a language to 1 iff it contains the empty word ε , and the transitions $\gamma_a : \varrho\bar{T} \rightarrow \varrho\bar{T}$ for $a \in X_0$ are given by $\gamma_a(L) = a^{-1}L$. Here $a^{-1}L = \{w \in X_0^* : aw \in L\}$ denotes the *left derivative* of L w.r.t. the letter a . Similarly, the *right derivatives* of L are defined by $La^{-1} = \{w \in X_0^* : wa \in L\}$ for $a \in X_0$.

The coalgebra $\varrho\bar{T}$ is characterized by a universal property: every finite \bar{T} -coalgebra has a unique coalgebra homomorphism into it (which sends a state to the language it accepts in the classical sense of automata theory). A finite \bar{T} -coalgebra is called a *subcoalgebra* of $\varrho\bar{T}$ if this unique morphism is injective, i.e. distinct states accept distinct languages. In [1] we related finite \bar{T} -coalgebras in \mathcal{C} to finite F -algebras in the predual category \mathcal{D} . Recall that $X = \Psi X_0$ implies $FA = I + X \otimes A \cong I + \coprod_{X_0} A$, so F -algebras $FA \xrightarrow{\delta} A$ can be represented as triples (A, δ_a, i) with $\delta_a : A \rightarrow A$ ($a \in X_0$) and $i : I \rightarrow A$. They correspond to automata in \mathcal{D} with inputs from the alphabet X_0 and without final states.

Proposition 6.4 (see [1]). (a) *The categories of finite \bar{T} -coalgebras and finite F -algebras are dually equivalent. The equivalence maps a finite \bar{T} -coalgebra $Q = (Q, \gamma_a, f)$ to its dual F -algebra $\hat{Q} = (\hat{Q}, \hat{\gamma}_a, \hat{f})$:*

$$(\bar{Y} \xleftarrow{f} Q \xrightarrow{\gamma_a} Q) \quad \mapsto \quad (I \xrightarrow{\hat{f}} \hat{Q} \xleftarrow{\hat{\gamma}_a} \hat{Q}).$$

(b) A finite \bar{T} -coalgebra Q is a subcoalgebra of $\varrho\bar{T}$ iff its dual F -algebra \hat{Q} is a quotient of the initial F -algebra X^\otimes .

Example 6.5. For a finite \bar{T} -coalgebra (Q, γ_a, f) in \mathbf{BA} the dual F -algebra \hat{Q} has as states the atoms of Q , and the initial state is the unique atomic final state of Q . Moreover, there is a transition $z \xrightarrow{a} z'$ for $a \in X_0$ in \hat{Q} iff z' is the unique atom with $\gamma_a(z') \geq z$ in Q .

By a *local variety of languages over X_0* in \mathcal{C} we mean a subcoalgebra V of $\varrho\bar{T}$ closed under right derivatives (i.e. $L \in |V|$ implies $La^{-1} \in |V|$ for all $a \in X_0$). Note that a local variety is also closed under the \mathcal{C} -algebraic operations of $\varrho\bar{T}$, being a subalgebra of $\varrho\bar{T}$ in \mathcal{C} , and under left derivatives, being a subcoalgebra of $\varrho\bar{T}$.

Example 6.6. A local variety of languages in \mathbf{BA} is a set of regular languages over X_0 closed under the boolean operations (union, intersection and complement) as well as left and right derivatives. This concept was introduced by Gehrke, Grigorieff and Pin [11].

For the following proposition recall that from every X_0 -generated \mathcal{D} -monoid one can derive an F -algebra, see Definition 2.19.

Proposition 6.7 (see [1]). *A finite subcoalgebra V of $\varrho\bar{T}$ is a local variety iff its dual F -algebra \hat{V} is derived from an X_0 -generated \mathcal{D} -monoid.*

In other words, given a finite local variety $V \hookrightarrow \varrho\bar{T}$ in \mathcal{C} , there exists a unique \mathcal{D} -monoid structure on \hat{V} making the unique (surjective) F -algebra homomorphism $e_{\hat{V}} : X^\otimes \rightarrow \hat{V}$ a \mathcal{D} -monoid morphism. Hence the monoid multiplication on \hat{V} is (well-)defined by $e_{\hat{V}}(x) \bullet e_{\hat{V}}(y) := e_{\hat{V}}(x \bullet y)$ for all $x, y \in X^\otimes$, and the unit is the initial state of the F -algebra \hat{V} .

Remark 6.8. A *pointed \bar{T} -coalgebra* is a \bar{T} -coalgebra (Q, γ_a, f) equipped with an initial state $i : \bar{I} \rightarrow Q$. Observe that every finite pointed \bar{T} -coalgebra (Q, γ_a, f, i) dualizes to a finite \mathcal{D} -automaton $(\hat{Q}, \hat{\gamma}_a, \hat{f}, \hat{i})$. The *language* of (Q, δ_a, f, i) is the function


$$L_Q : X_0^* \rightarrow |\bar{Y}|, \quad a_1 \cdots a_n \mapsto f \cdot \delta_{a_n} \cdots \delta_{a_1} \cdot i,$$

where $|\bar{Y}|$ is identified with $|Y|$ as in (6.1). Letting $m_Q : Q \rightarrow \varrho\bar{T}$ denote the unique coalgebra homomorphism, L_Q is precisely the element of $\varrho\bar{T}$ determined by $I \xrightarrow{i} Q \xrightarrow{m_Q} \varrho\bar{T}$. Since $|\bar{Y}| = |Y|$ and $X^\otimes = \Psi X_0^*$, the function $L_Q : X_0^* \rightarrow |\bar{Y}|$ can be identified with its adjoint transpose $L_Q^\circ : X^\otimes \rightarrow Y$, i.e. with a language in \mathcal{D} . The *reversal* of a language $L : X^\otimes \rightarrow Y$ in \mathcal{D} is $L^{\text{rev}} = L \cdot \text{rev} : X^\otimes \rightarrow Y$, where $\text{rev} : X^\otimes \rightarrow X^\otimes$ denotes the unique morphism of \mathcal{D} extending the function $X_0^* \rightarrow X_0^*$ that reverses words.

Proposition 6.9 (see [3]). *The language accepted by a finite pointed \bar{T} -coalgebra is the reversal of the language accepted by its dual \mathcal{D} -automaton.*

If a finite X_0 -generated \mathcal{D} -monoid $e : X^\otimes \rightarrow M$ recognizes a language $L : X^\otimes \rightarrow Y$ via $f : M \rightarrow Y$, i.e. $L = (X^\otimes \xrightarrow{e} M \xrightarrow{f} Y)$, we dually get the morphism $\bar{I} \xrightarrow{i} V \xrightarrow{m} \varrho\bar{T}$ (where V is the local variety dual to M , i is the dual morphism of f and m is the unique coalgebra homomorphism) choosing the element L^{rev} of $\varrho\bar{T}$. Now suppose that L is a regular language, and let V_L be the finite local variety of languages dual to the syntactic \mathcal{D} -monoid $\text{Syn}(L)$, see Proposition 6.7. The universal property of $\text{Syn}(L)$ in Definition 3.6 then dualizes as follows: (i) V_L is a local variety containing L^{rev} , and (ii) for every local variety $V \hookrightarrow \varrho\bar{T}$

containing L^{rev} , the local variety V_L is contained in V . In other words, V_L is the *smallest* local variety containing L^{rev} .



In summary, we have proved the following dual characterization of syntactic \mathcal{D} -monoids:

Theorem 6.10. *For every regular language L the syntactic \mathcal{D} -monoid $\text{Syn}(L)$ is dual to the smallest local variety of languages over X_0 in \mathcal{C} containing L^{rev} .*

Example 6.11. For $\mathcal{C} = \mathbf{BA}$ and $\mathcal{D} = \mathbf{Set}$ the previous theorem gives the following construction of the syntactic monoid of a regular language $L \subseteq X^*$:

- (1) Form the smallest local variety of languages $V_L \subseteq \varrho\bar{T}$ containing L^{rev} . Hence V_L is the closure of the (finite) set of all both-sided derivatives $u^{-1}L^{\text{rev}}v^{-1} = \{w \in X^* : u w v \in L^{\text{rev}}\}$ ($u, v \in X^*$) under union, intersection and complement.
- (2) Compute the F -algebra \widehat{V}_L dual to the coalgebra V_L , which is a quotient $e_{\widehat{V}_L} : X^{\otimes} \rightarrow \widehat{V}_L$. The states of \widehat{V}_L are the atoms of V_L , and the initial state is the unique atom $i \in V_L$ containing the empty word. Given atoms $z, z' \in V_L$ and $a \in X$, there is a transition $z \xrightarrow{a} z'$ in \widehat{V}_L iff z' is the (unique) atom with $z \subseteq a^{-1}z'$.
- (3) Define a monoid multiplication on \widehat{V}_L as follows: given states $z, z' \in \widehat{V}_L$, choose words $w, w' \in X^*$ with $e_{\widehat{V}_L}(w) = z$ and $e_{\widehat{V}_L}(w') = z'$ (i.e. $i \xrightarrow{w} z$ and $i \xrightarrow{w'} z'$ in \widehat{V}_L). Then put $z \bullet z' = e_{\widehat{V}_L}(ww')$; this is the state reached on input ww' , i.e. $i \xrightarrow{ww'} z \bullet z'$. The resulting monoid (with multiplication \bullet and unit i) is $\text{Syn}(L)$.

By dropping right derivatives and using the correspondence between finite subcoalgebras of $\varrho\bar{T}$ and finite quotient algebras of X^{\otimes} , one also gets the following dual characterization of minimal \mathcal{D} -automata; cf. also [6] for a closely related dual prespective on minimal automata.

Theorem 6.12. *For every regular language L the minimal \mathcal{D} -automaton for L is dual to the smallest subcoalgebra of $\varrho\bar{T}$ containing L^{rev}*

Remark 6.13. Our assumption that Y and \bar{Y} have two elements is inessential. Without this assumption, the rational coalgebra $\varrho\bar{T}$ is not carried by the set of regular languages, but more generally by the set of *regular behaviors*, i.e. functions $b : X_0^* \rightarrow |Y|$ realized by finite Moore automata with output set $|Y|$. The coalgebra structure is given by the output map $b \mapsto b(\varepsilon)$, and transitions $b \xrightarrow{a} a^{-1}b$ for $a \in X_0$, where $a^{-1}b$ is the (generalized) left derivative of b defined by $a^{-1}b(w) = b(aw)$. (Right derivatives are defined analogously.) A *local variety of behaviors over X_0* in \mathcal{C} is again a subcoalgebra of $\varrho\bar{T}$ closed under right derivatives. All results of this section hold for this more general setting, see Section 5 of [3] for details. In particular, this allows us to cover the case $\mathcal{C} = \mathcal{D} = \mathbf{Vec}(\mathbb{K})$ for an arbitrary finite field \mathbb{K} . In this case Theorem 6.10 states that the syntactic associative algebra of a rational weighted language $L : X_0^* \rightarrow \mathbb{K}$ dualizes to the smallest set of rational weighted languages that contains L^{rev} and is closed under scalar multiplication, addition and left and right derivatives.

7. CONCLUSIONS AND FUTURE WORK

We proposed the first steps of a categorical theory of algebraic language recognition. Despite our assumption that \mathcal{D} be a commutative variety, we have seen that a number of our definitions, constructions and proofs work in any symmetric monoidal closed category with enough structure. The next step should be the treatment of algebraic recognition beyond languages of finite words, including such examples as Wilke algebras [22] (representing ω -languages) or forest algebras [9] (representing tree languages). A first categorical approach to such structures appears in the work of Bojanczyk [8] who works with monads on sorted sets rather than monoids in a variety \mathcal{D} . We expect that a unification of these two directions is possible.

One of the leading themes of algebraic automata theory is the classification of languages in terms of their syntactic algebras. For instance, by Schutzenberger’s theorem a language is star-free iff its syntactic monoid is aperiodic. We hope that our conceptual view of syntactic monoids (notably their dual characterization in Theorem 6.10) can contribute to a duality-based approach to such results.

REFERENCES

- [1] Jir Adamek, Stefan Milius, Robert S. R. Myers, and Henning Urbat. Generalized Eilenberg Theorem I: Local Varieties of Languages. In Anca Muscholl, editor, *Proc. Foundations of Software Science and Computation Structures (FoSSaCS)*, volume 8412 of *Lecture Notes Comput. Sci.*, pages 366–380. Springer, 2014.
- [2] Jir Adamek, Stefan Milius, and Henning Urbat. Syntactic monoids in a category. In *Proc. 6th Conference on Algebra and Coalgebra in Computer Science (CALCO’15)*, Leibniz International Proceedings in Informatics (LIPIcs), 2015.
- [3] Jir Adamek, Robert S. R. Myers, Henning Urbat, and Stefan Milius. Varieties of languages in a category. In *30th Annual ACM/IEEE Symposium on Logic in Computer Science*. IEEE, 2015.
- [4] Adolfo Ballester-Bolinches, Enric Cosme-Llopez, and Jan J. M. M. Rutten. The dual equivalence of equations and coequations for automata. *Inform. and Comput.*, 244:49–75, 2014.
- [5] Bernhard Banaschewski and Evelyn Nelson. Tensor products and bimorphisms. *Canad. Math. Bull.*, 19:385–402, 1976.
- [6] Nick Bezhanishvili, Clemens Kupke, and Prakash Panangaden. Minimization via duality. In Luke Ong and Ruy de Queiroz, editors, *Logic, Language, Information and Computation*, volume 7456 of *Lecture Notes in Computer Science*, pages 191–205. Springer Berlin Heidelberg, 2012.
- [7] Stephen L. Bloom. Varieties of ordered algebras. *J. Comput. Syst. Sci.*, 13(2):200–212, 1976.
- [8] Mikoaj Bojanczyk. Recognisable languages over monads. In Igor Potapov, editor, *Developments in Language Theory*, volume 9168 of *LNCS*, pages 1–13. Springer, 2015. Full version: <http://arxiv.org/abs/1502.04898>.
- [9] Mikoaj Bojanczyk and Igor Walukiewicz. Forest algebras. In *Automata and Logic: History and Perspectives*, pages 107–132, 2006.
- [10] Marcello M. Bonsangue, Stefan Milius, and Alexandra Silva. Sound and complete axiomatizations of coalgebraic language equivalence. *ACM Trans. Comput. Log.*, 14(1:7), 2013.
- [11] Mai Gehrke, Serge Grigorieff, and Jean-eric Pin. Duality and equational theory of regular languages. In *Proc. ICALP 2008, Part II*, volume 5126 of *Lecture Notes Comput. Sci.*, pages 246–257. Springer, 2008.
- [12] Joseph A. Goguen. Discrete-time machines in closed monoidal categories. I. *J. Comput. Syst. Sci.*, 10(1):1–43, 1975.
- [13] Anders Kock. Monads on symmetric monoidal closed categories. *Arch. Math.*, 21:1–10, 1970.
- [14] Saunders Mac Lane. *Categories for the working mathematician*. Springer, 2nd edition, 1998.
- [15] Jean-eric Pin. Mathematical foundations of automata theory. Available at <http://www.liafa.jussieu.fr/~jep/PDF/MPRI/MPRI.pdf>.
- [16] Jean-eric Pin. A variety theorem without complementation. *Russian Mathematics*, 39:80–90, 1995.

- [17] Jean-Éric Pin and Pascal Weil. A Reiterman theorem for pseudovarieties of finite first-order structures. *Alg. Univ.*, 35:577–595, 1996.
- [18] Libor Polák. Syntactic semiring of a language. In Jiří Sgall, Aleš Pultr, and Petr Kolman, editors, *Proc. International Symposium on Mathematical Foundations of Computer Science (MFCS)*, volume 2136 of *Lecture Notes Comput. Sci.*, pages 611–620. Springer, 2001.
- [19] Michael O. Rabin and Dana S. Scott. Finite automata and their decision problems. *IBM J. Res. Dev.*, 3(2):114–125, April 1959.
- [20] Christophe Reutenauer. Séries formelles et algèbres syntactiques. *J. Algebra*, 66:448–483, 1980.
- [21] Jan J. M. M. Rutten. Universal coalgebra: a theory of systems. *Theoret. Comput. Sci.*, 249(1):3–80, 2000.
- [22] Thomas Wilke. An Eilenberg Theorem for Infinity-Languages. In *Proc. ICALP 91*, pages 588–599. Springer, 1991.